

# Uniform confidence bands in deconvolution with unknown error distribution

Kengo Kato\*

Graduate School of Economics, University of Tokyo

and

Yuya Sasaki

Department of Economics, Johns Hopkins University

January 31, 2017

## Abstract

This paper develops a method to construct uniform confidence bands in deconvolution when the error distribution is unknown. We work with the setting where an auxiliary sample from the error distribution is available and the error density is ordinary smooth. The construction is based upon the “intermediate” Gaussian approximation and the Gaussian multiplier bootstrap, but not on explicit limit distributions such as Gumbel distributions, which enables us to prove validity of the proposed multiplier bootstrap confidence band under mild regularity conditions. Importantly, the previous literature on this topic has focused on the case where the error distribution is known, but has not formally covered the case with unknown error distribution. However, in many applications, the assumption that the error density is known is unrealistic, and the present paper fills this important void. In addition, we also discuss extensions of the results on confidence bands to super-smooth error densities. We conduct simulation studies to verify the performance of the multiplier bootstrap confidence band in the finite sample. Finally, we apply our method to the Outer Continental Shelf (OCS) Auction Data and draw confidence bands for the density of common values of mineral rights on oil and gas tracts.

---

\*K. Kato is supported by Grant-in-Aid for Scientific Research (C) (15K03392) from the JSPS.

*Keywords:* bootstrap, confidence band, deconvolution, empirical process theory.

# 1 Introduction

Consider the deconvolution model

$$Y = X + \varepsilon, \tag{1.1}$$

where  $X$  and  $\varepsilon$  are independent, real-valued random variables with unknown densities  $f_X$  and  $f_\varepsilon$ , respectively. We observe  $Y$ , but do not observe  $X$  nor  $\varepsilon$ . The variable  $\varepsilon$  is referred to as a measurement error. The model (1.1) implies that the density  $f_Y$  of  $Y$  exists and is given by the convolution of  $f_X$  and  $f_\varepsilon$ , namely,

$$f_Y(y) = (f_X * f_\varepsilon)(y) = \int_{\mathbb{R}} f_X(x) f_\varepsilon(y - x) dx. \tag{1.2}$$

The deconvolution problem refers to the problem of estimating and making inference on  $f_X$  from available data.

The goal of this paper is to develop a method to construct uniform confidence bands for  $f_X$  when the error density  $f_\varepsilon$  is unknown. To deal with unknown error distribution, in addition to the availability of an independent sample  $Y_1, \dots, Y_n$  from  $f_Y$ , we assume that an auxiliary independent sample  $\eta_1, \dots, \eta_m$  from  $f_\varepsilon$  is available where  $m = m_n \rightarrow \infty$  as  $n \rightarrow \infty$ . This assumption is satisfied in various ways depending on an application of interest. For example, in natural science, measurement errors are often due to measuring devices; in such cases, one can obtain preliminary calibration measures in the absence of signal, which produce a sample from the error distribution; see, for example, the introduction of Comte and Lacour (2011) for such cases. Another case, which accommodates a wider spectrum of relevant applications, is where panel data or repeated measurements  $(Y^{(1)}, Y^{(2)})$  for a common signal  $X$  with errors  $(\varepsilon^{(1)}, \varepsilon^{(2)})$  such that the conditional distribution of the one given another is symmetric are available (see Example 2.1 ahead). Under these two representative situations with unknown  $f_\varepsilon$ , we develop a deconvolution estimator and

confidence bands for  $f_X$  based on a multiplier bootstrap. We use a standard deconvolution kernel density estimator that goes back to Carroll and Hall (1988) and Stefanski and Carroll (1990), but replace the error characteristic function by the empirical characteristic function based on the auxiliary sample  $\eta_1, \dots, \eta_m$  from  $f_\varepsilon$ .

It is well understood that the difficulty of estimating  $f_X$  critically depends on how fast the modulus of the error characteristic function  $\varphi_\varepsilon(t) = \mathbb{E}[e^{it\varepsilon}]$  with  $i = \sqrt{-1}$  decays as  $|t| \rightarrow \infty$ , in addition to the smoothness of  $f_X$  (the faster  $|\varphi_\varepsilon(t)|$  decays as  $|t| \rightarrow \infty$ , the more difficult estimation of  $f_X$  will be). Informally, the error density  $f_\varepsilon$  is said to be *ordinary smooth* if  $|\varphi_\varepsilon(t)|$  decays at most polynomially fast as  $|t| \rightarrow \infty$ , while  $f_\varepsilon$  is said to be *super-smooth* if  $|\varphi_\varepsilon(t)|$  decays exponentially fast as  $|t| \rightarrow \infty$  (cf. Fan, 1991a). We first consider ordinary smooth error densities and prove asymptotic validity of the multiplier bootstrap confidence band under mild regularity conditions. In this ordinary smooth case, the sample size  $m$  for  $f_\varepsilon$  need not be large in comparison with  $n$ . As a corollary, we derive uniform convergence rates of the deconvolution kernel density estimator with the estimated error characteristic function. Furthermore, we extend the results on confidence bands to super-smooth error densities; in the super-smooth case, however, we require  $m/n \rightarrow \infty$  for a technical reason.

We conduct simulation studies to verify the performance of the multiplier bootstrap confidence band in the finite sample. The simulation studies show that the simulated coverage probabilities are very close to nominal coverage probabilities even with sample sizes as small as 250 and 500, suggesting practical benefits of our confidence band. Finally, following Li et al. (2000), we apply our method to the Outer Continental Shelf (OCS) Auction Data (Hendricks et al., 1987) and draw confidence bands for the density of common values of mineral rights on oil and gas tracts in the Gulf of Mexico. In the empirical auction literature, obtaining confidence intervals/bands for a deconvolution density with unknown

error distribution is of interest (e.g., Krasnokutskaya, 2011), and practitioners have implemented nonparametric bootstrap without a theoretical support for its validity. We draw valid confidence bands for a deconvolution density, and provide statistical support for some qualitative features of the common value density that Li et al. (2000) find visually in their estimate. In order to illustrate the advantage of our method over existing alternatives which assume to know the error density, we also construct a uniform confidence band by assuming that the error density is known to take a particular form. We find qualitative differences between the two confidence bands, highlighting practical values of our approach which requires no prior knowledge of the error distribution.

The literature related to this paper is broad. We refer to Carroll et al. (2006), Meister (2009), and Horowitz (2009) as general references on measurement error models and deconvolution. In the present paper, among other approaches, we build upon the deconvolution kernel density estimation method, which is pioneered by Carroll and Hall (1988); Stefanski and Carroll (1990); Fan (1991a,b). These earlier studies, as well as many of more recent papers, focus on the case where the error density  $f_\varepsilon$  is assumed to be known, even though it is often unknown in applications.

The deconvolution problem with unknown  $f_\varepsilon$  is studied by Diggle and Hall (1993); Efromovich (1997); Neumann (1997); Johannes (2009); Comte and Lacour (2011); Dattner et al. (2016). These papers assume the availability of auxiliary measurements from the error distribution. Horowitz and Markatou (1996) and Li and Vuong (1998) consider to estimate a deconvolution density from repeated measurements (panel data) of  $Y$ , instead of assuming measurements from the error distribution *per se*; see also Neumann (2007), Delaigle et al. (2008), and Bohnomme and Robin (2010) for further developments. In addition, a recent work by Delaigle and Hall (2016) relaxes the requirement of repeated measurements under the assumption of a symmetric error distribution. Despite the richness of this literature,

however, uniform confidence bands for  $f_X$  have not been developed in any of these papers allowing for unknown  $f_\varepsilon$ .

The deconvolution problem is a statistical ill-posed inverse problem, and developing formal theories for inference in ill-posed inverse problems tends to be challenging. In fact, it is only (relatively) recently that studies on uniform confidence bands in deconvolution have appeared, and those previous studies focus on the case where  $f_\varepsilon$  is known. To the best of our knowledge, Bissantz et al. (2007) is the first paper that formally studies uniform confidence bands in deconvolution. They assume that  $f_\varepsilon$  is known and ordinary smooth, and prove, under a number of technical conditions, a Smirnov-Bickel-Rosenblatt type limit theorem (cf. Smirnov, 1950; Bickel and Rosenblatt, 1973) for the deconvolution kernel density estimator based on the Komlós-Major-Tusnády (KMT) strong approximation (Komlós et al., 1975) and extreme value theory (cf. Leadbetter et al., 1983). Namely, they prove that the supremum deviation of the deconvolution kernel density estimator, suitably normalized, converges in distribution to a Gumbel distribution. Further, they prove consistency of the nonparametric bootstrap. See also Bissantz and Holzmann (2008). Interestingly, when the error density is super-smooth, van Es and Gugushvili (2008) show that the limit distribution of the supremum deviation of the deconvolution kernel density estimator in general differs from Gumbel distributions. We also refer to Lounici and Nickl (2011); Schmidt-Hieber et al. (2013); Delaigle et al. (2015).

Importantly, none of these papers formally studies the case where the error density  $f_\varepsilon$  is unknown. It should be noted that Delaigle et al. (2015, Section 4.2) discuss how to possibly accommodate the case of unknown error density, but a theory to support this argument is not provided.<sup>1</sup> While the effect of pre-estimating the unknown error characteristic function

---

<sup>1</sup>The focus in Delaigle et al. (2015) is on pointwise confidence intervals for nonparametric regression functions, and differs from our objective to conduct uniform inference on nonparametric density functions.

for the purpose of estimating  $f_X$  is modest, its effect for the purpose of inference on  $f_X$  is non-trivial. We attempt to contribute to the literature on deconvolution by formally establishing a method to construct uniform confidence bands for  $f_X$  under the realistic situation where the error density  $f_\varepsilon$  is unknown. It is worth pointing out that the multiplier bootstrap confidence band proposed in the present paper is valid for both cases where the error density is ordinary- and super-smooth (although in the latter case we require  $m/n \rightarrow \infty$ ), despite the fact that, as already mentioned, the limit distributions of the supremum deviation of the deconvolution kernel density estimator (with known  $f_\varepsilon$ ) in general differ in those two cases.

From a technical point of view, the present paper builds upon non-trivial applications of the “intermediate” Gaussian and multiplier bootstrap approximation theorems developed in Chernozhukov et al. (2014a,b, 2016). These approximation theorems are applicable to the general empirical process under weaker regularity conditions than those for the KMT and Gumbel approximations. In particular, they enable us to prove validity of the multiplier bootstrap confidence band without relying on explicit limit distributions such as Gumbel distributions and thereby under mild regularity conditions. See the discussion in Remark 3.4 in Section 3. However, we stress that those theorems are not directly applicable to our problems and substantial work is needed to derive our results. This is because: 1) the “deconvolution” kernel  $K_n$  (see Section 2 ahead) is implicitly defined via the Fourier inversion, and verifying conditions in those approximation theorems with the deconvolution kernel  $K_n$  is non-trivial; 2) the error density  $f_\varepsilon$  is unknown and we have to work with the estimated deconvolution kernel  $\widehat{K}_n$ , and so the estimation error has to be taken into account, which requires delicate cares since  $\widehat{K}_n$  depends on the inverse of the estimated error characteristic function, and the error characteristic function is, although it is assumed to be non-vanishing on the entire real line, approaching zero in the tail; see the

discussion after Theorem 3.1.

The rest of the paper is organized as follows. Section 2 presents our methodology of constructing confidence bands for  $f_X$ . Section 3 presents the main theoretical results of this paper where we consider the ordinary smooth case. Section 4 presents the results on numerical simulations. Section 5 presents an empirical application of our method. Section 6 presents extensions of the results on confidence bands to super-smooth error densities. The supplementary material contains all the proofs, additional simulation results, and a description on the empirical data.

## 1.1 Notation

For  $a, b \in \mathbb{R}$ , let  $a \wedge b = \min\{a, b\}$ . For a non-empty set  $T$  and a (complex-valued) function  $f$  on  $T$ , we use the notation  $\|f\|_T = \sup_{t \in T} |f(t)|$ . Let  $\ell^\infty(T)$  denote the Banach space of all bounded real-valued functions on  $T$  with norm  $\|\cdot\|_T$ . The Fourier transform of an integrable function  $f$  on  $\mathbb{R}$  is defined by  $\varphi_f(t) = \int_{\mathbb{R}} e^{itx} f(x) dx$  for  $t \in \mathbb{R}$ , where  $i = \sqrt{-1}$  denotes the imaginary unit throughout the paper. We refer to Folland (1999) as a basic reference on Fourier analysis.

## 2 Methodology

In this section, we informally present our methodology to construct confidence bands for  $f_X$ . The formal analysis of our methodology will be carried out in the following sections.

Let  $\varphi_Y, \varphi_X$ , and  $\varphi_\varepsilon$  denote the Fourier transforms (the characteristic functions) of  $f_Y, f_X$ , and  $f_\varepsilon$ , respectively. The model (1.1) implies that these characteristic functions satisfy the relation

$$\varphi_Y(t) = \varphi_X(t)\varphi_\varepsilon(t) \quad \text{for all } t \in \mathbb{R}. \quad (2.1)$$

Provided that  $\varphi_\varepsilon$  does not vanish on  $\mathbb{R}$  and  $|\varphi_X|$  is integrable on  $\mathbb{R}$ , the Fourier inversion formula yields that  $f_X$  is recovered from  $f_Y$  and  $f_\varepsilon$  via

$$f_X(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \frac{\varphi_Y(t)}{\varphi_\varepsilon(t)} dt. \quad (2.2)$$

Precisely speaking, if  $|\varphi_X|$  is integrable on  $\mathbb{R}$ , then  $f_X$  has a version that is bounded and continuous, and satisfies (2.2) for each  $x \in \mathbb{R}$ ; we take this version in what follows. Suppose that independent copies  $Y_1, \dots, Y_n$  of  $Y$  are observed. To describe an approach to constructing confidence bands for  $f_X$ , suppose first that the distribution of  $\varepsilon$  were known. In that case, a standard kernel estimator of  $f_X$  is given by

$$\hat{f}_X^*(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \hat{\varphi}_Y(t) \frac{\varphi_K(th_n)}{\varphi_\varepsilon(t)} dt = \frac{1}{nh_n} \sum_{j=1}^n K_n((x - Y_j)/h_n),$$

where

$$\hat{\varphi}_Y(t) = \frac{1}{n} \sum_{j=1}^n e^{itY_j} \quad \text{and} \quad K_n(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \frac{\varphi_K(t)}{\varphi_\varepsilon(t/h_n)} dt,$$

and where  $K : \mathbb{R} \rightarrow \mathbb{R}$  is a real-valued kernel function (i.e., an integrable function that integrates to one) such that its Fourier transform  $\varphi_K$  is supported in  $[-1, 1]$  (i.e.,  $\varphi_K(t) = 0$  for all  $|t| > 1$ ), and  $h_n$  is a sequence of positive numbers (bandwidths) such that  $h_n \rightarrow 0$  (cf. Carroll and Hall, 1988; Stefanski and Carroll, 1990). Note that the function  $K_n$  is real-valued, and bounded by compactness of the support of  $\varphi_K$ . The function  $K_n$  is called a *deconvolution kernel*.

Suppose that we are interested in constructing a confidence band for  $f_X$  on a compact interval  $I \subset \mathbb{R}$ . A confidence band  $\mathcal{C}_n$  at level  $(1 - \tau)$ , where  $\tau \in (0, 1)$  is given, is a family of random intervals  $\mathcal{C}_n = \{\mathcal{C}_n(x) = [c_L(x), c_U(x)] : x \in I\}$  such that

$$\mathbb{P}\{f_X(x) \in [c_L(x), c_U(x)] \forall x \in I\} \geq 1 - \tau.$$

An approach to constructing such a confidence band is based on approximating the distribution of the supremum in absolute value of the following stochastic process:

$$Z_n^*(x) = \frac{\widehat{f}_X^*(x) - \mathbb{E}[\widehat{f}_X^*(x)]}{\sqrt{\text{Var}(\widehat{f}_X^*(x))}} = \frac{\sqrt{nh_n}\{\widehat{f}_X^*(x) - \mathbb{E}[\widehat{f}_X^*(x)]\}}{\sigma_n(x)}, \quad x \in I,$$

where we assume that  $\sigma_n^2(x) = \text{Var}(K_n((x - Y)/h_n)) > 0$  for all  $x \in I$ , and  $\sigma_n(x)$  is defined by  $\sigma_n(x) = \sqrt{\sigma_n^2(x)}$ . By setting

$$c_n^*(1 - \tau) = (1 - \tau)\text{-quantile of } \|Z_n^*\|_I,$$

we see that a band of the form

$$\mathcal{C}_n^*(x) = \left[ \widehat{f}_X^*(x) - \frac{\sigma_n(x)}{\sqrt{nh_n}} c_n^*(1 - \tau), \widehat{f}_X^*(x) + \frac{\sigma_n(x)}{\sqrt{nh_n}} c_n^*(1 - \tau) \right], \quad x \in I \quad (2.3)$$

contains  $\{\mathbb{E}[\widehat{f}_X^*(x)] : x \in I\}$  with probability at least  $(1 - \tau)$ , since

$$\mathbb{P}\{\mathbb{E}[\widehat{f}_X^*(x)] \in \mathcal{C}_n^*(x) \forall x \in I\} = \mathbb{P}\{\|Z_n^*\|_I \leq c_n^*(1 - \tau)\} \geq 1 - \tau.$$

Provided that the bias  $\|f_X(\cdot) - \mathbb{E}[\widehat{f}_X^*(\cdot)]\|_I$  is sufficiently small (which is typically achieved by choosing undersmoothing bandwidths), the band of the form (2.3) is a valid confidence band for  $f_X$  on  $I$  at level approximately  $(1 - \tau)$ .

Constructing a band of the form (2.3) is, however, infeasible since both the distribution of  $\|Z_n^*\|_I$  and the variance function  $\sigma_n^2(\cdot)$  are unknown. More fundamentally, in many applications, the error density  $f_\varepsilon$  (or more directly the error characteristic function  $\varphi_\varepsilon$ ) is unknown, and so the deconvolution kernel estimator  $\widehat{f}_X^*$  is simply infeasible. In this paper, we assume that  $f_\varepsilon$  is unknown, but assume the availability of an independent sample  $\eta_1, \dots, \eta_m$  from  $f_\varepsilon$  where  $m = m_n \rightarrow \infty$  as  $n \rightarrow \infty$ . We do not assume that  $(\eta_1, \dots, \eta_m)$  are independent from  $(Y_1, \dots, Y_n)$ . A simple such case is where auxiliary experiments provide a

sample  $\eta_1, \dots, \eta_m$ , which is typically the case when measurement error is due to measuring devices. Another typical example where such observations are available is the case where we observe repeated measurements on  $X$  with errors such that the conditional distribution of the one given another is symmetric.

**Example 2.1** (Carroll et al. (2006), p.298). Suppose that we observe repeated measurements on  $X$  with errors as  $Y^{(j)} = X + \varepsilon^{(j)}$  for  $j = 1, 2$  where  $X$  and  $(\varepsilon^{(1)}, \varepsilon^{(2)})$  are independent ( $\varepsilon^{(1)}$  and  $\varepsilon^{(2)}$  need not to be independent, nor do they have common distribution). Then we have  $\underbrace{(Y^{(1)} + Y^{(2)})/2}_{=:Y} = X + \underbrace{(\varepsilon^{(1)} + \varepsilon^{(2)})/2}_{=: \varepsilon}$  and  $(\varepsilon^{(1)} - \varepsilon^{(2)})/2 = (Y^{(1)} - Y^{(2)})/2$ . So provided that  $\varepsilon^{(1)} + \varepsilon^{(2)}$  has the same distribution as  $\varepsilon^{(1)} - \varepsilon^{(2)}$ , which is true if the conditional distribution of  $\varepsilon^{(2)}$  given  $\varepsilon^{(1)}$  is symmetric,  $\eta := (Y^{(1)} - Y^{(2)})/2$  has the same distribution as  $\varepsilon = (\varepsilon^{(1)} + \varepsilon^{(2)})/2$ . In this example,  $m = n$ .

In such a case, a natural estimator of  $\varphi_\varepsilon$  is the empirical characteristic function based on  $\eta_1, \dots, \eta_m$ , namely,

$$\widehat{\varphi}_\varepsilon(t) = \frac{1}{m} \sum_{j=1}^m e^{it\eta_j}.$$

Suppose for a while that  $\inf_{|t| \leq h_n^{-1}} |\widehat{\varphi}_\varepsilon(t)| > 0$  with probability approaching one (which is indeed guaranteed under the assumptions stated below); then we may estimate the deconvolution kernel  $K_n$  as

$$\widehat{K}_n(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \frac{\varphi_K(t)}{\widehat{\varphi}_\varepsilon(t/h_n)} dt. \quad (2.4)$$

Now, define the feasible version of  $\widehat{f}_X^*$  as

$$\widehat{f}_X(x) = \frac{1}{nh_n} \sum_{j=1}^n \widehat{K}_n((x - Y_j)/h_n).$$

(Formally think of  $\widehat{K}_n(x) = 0$  if the integral in (2.4) is not well-defined.) This estimator was first considered by Diggle and Hall (1993). Note that  $\widehat{K}_n$  is also real-valued. In addition, we may estimate the variance function  $\sigma_n^2(x)$  by

$$\widehat{\sigma}_n^2(x) = \frac{1}{n} \sum_{j=1}^n \widehat{K}_n^2((x - Y_j)/h_n) - \left( \frac{1}{n} \sum_{j=1}^n \widehat{K}_n((x - Y_j)/h_n) \right)^2.$$

Consider the stochastic process

$$\widehat{Z}_n(x) = \frac{\sqrt{nh_n}\{f_X(x) - \mathbb{E}[f_X^*(x)]\}}{\widehat{\sigma}_n(x)}, \quad x \in I,$$

where  $\widehat{\sigma}_n(x) = \sqrt{\widehat{\sigma}_n^2(x)}$ , and let

$$c_n(1 - \tau) = (1 - \tau)\text{-quantile of } \|\widehat{Z}_n\|_I.$$

(Formally think of  $\widehat{Z}_n(x)$  as 0 if  $\widehat{\sigma}_n(x) = 0$ ; the probability that  $\widehat{\sigma}_n(x) = 0$  for some  $x \in I$  is approaching zero under the assumptions stated below. The same rule applies to  $\widehat{Z}_n^\xi$  defined below.) Then a band of the form

$$\left[ \widehat{f}_X(x) - \frac{\widehat{\sigma}_n(x)}{\sqrt{nh_n}} c_n(1 - \tau), \widehat{f}_X(x) + \frac{\widehat{\sigma}_n(x)}{\sqrt{nh_n}} c_n(1 - \tau) \right], \quad x \in I$$

will be a valid confidence band for  $f_X$  on  $I$  at level approximately  $(1 - \tau)$  provided that the bias  $\|f_X(\cdot) - \mathbb{E}[f_X^*(\cdot)]\|_I$  is sufficiently small.

The quantiles of  $\|\widehat{Z}_n\|_I$  are still unknown, but it will be shown below that under suitable regularity conditions the distribution of  $\|\widehat{Z}_n\|_I$  can be approximated by that of the supremum in absolute value of a tight Gaussian random variable  $Z_n^G$  in  $\ell^\infty(I)$  with mean zero and the same covariance function as  $Z_n^*$ . So, we propose to estimate the quantiles of  $\|\widehat{Z}_n\|_I$  via the *Gaussian multiplier bootstrap* as in Chernozhukov et al. (2014b), described as follows.

Generate independent standard normal random variables  $\xi_1, \dots, \xi_n \sim N(0, 1)$ , independent of the data  $\mathcal{D}_n = \{Y_1, \dots, Y_n, \eta_1, \dots, \eta_m\}$ , and consider the stochastic process

$$\widehat{Z}_n^\xi(x) = \frac{1}{\widehat{\sigma}_n(x)\sqrt{n}} \sum_{j=1}^n \xi_j \left\{ \widehat{K}_n((x - Y_j)/h_n) - n^{-1} \sum_{j'=1}^n \widehat{K}_n((x - Y_{j'})/h_n) \right\}$$

for  $x \in I$ . Conditionally on the data  $\mathcal{D}_n$ ,  $\widehat{Z}_n^\xi(x), x \in I$  is a Gaussian process with mean zero and the covariance function “close” to that of  $Z_n^G$ . Hence we propose to estimate the quantile  $c_n(1 - \tau)$  by

$$\widehat{c}_n(1 - \tau) = \text{conditional } (1 - \tau)\text{-quantile of } \|\widehat{Z}_n^\xi\|_I \text{ given } \mathcal{D}_n,$$

and the resulting confidence band is

$$\widehat{\mathcal{C}}_n(x) = \left[ \widehat{f}_X(x) - \frac{\widehat{\sigma}_n(x)}{\sqrt{nh_n}} \widehat{c}_n(1 - \tau), \widehat{f}_X(x) + \frac{\widehat{\sigma}_n(x)}{\sqrt{nh_n}} \widehat{c}_n(1 - \tau) \right], \quad x \in I. \quad (2.5)$$

How to choose the bandwidth is an important yet difficult problem in any nonparametric inference. Practical choice of the bandwidth will be discussed in Section 4.2.

Note that the analysis below covers the case where  $I$  is singleton, i.e.,  $I = \{x_0\}$ , and in that case, the above confidence band gives a confidence interval for  $f_X(x_0)$ .

In practice, the presence of  $\widehat{\varphi}_\varepsilon$  in the denominator in the integrand in (2.4) could make the estimate  $\widehat{f}_X$  numerically unstable; an approach to cope with this problem is to restrict the integral in (2.4) to the set  $\{t \in \mathbb{R} : |\widehat{\varphi}_\varepsilon(t)| \geq m^{-1/2}\}$  (cf. Neumann, 1997). Likewise, in practice, replacing  $\widehat{\sigma}_n(x)$  by  $\max\{\widehat{\sigma}_n(x), \sqrt{h_n}\}$  in the definition of  $\widehat{Z}_n$  and  $\widehat{Z}_n^\xi$  would make resulting confidence bands numerically more stable. These modifications do not alter the asymptotic results presented below, and we will work with the original definitions of  $\widehat{K}_n$  and  $\widehat{\sigma}_n$ .

**Remark 2.1** (Independence between  $X$  and  $\varepsilon$ ). In the present paper, we work with the classical measurement error setting, namely, we assume that  $X$  and  $\varepsilon$  are independent.

However, for our theoretical results to hold, full independence between  $X$  and  $\varepsilon$  is not necessary. Instead, we only require condition (2.1), which can hold even when  $X$  and  $\varepsilon$  are not independent. Schennach (2013) argues that condition (2.1) is “as weak as a conditional mean assumption”.

### 3 Main results

In this section, we present the theorems that provide asymptotic validity of our confidence band described in the previous section. We first consider the case where the error density  $f_\varepsilon$  is ordinary smooth. We begin with stating the assumptions that guarantee our asymptotic results.

**Assumption 1.** *The function  $|\varphi_X|$  is integrable on  $\mathbb{R}$ .*

**Assumption 2.** *Let  $K$  be a real-valued integrable function (kernel) on  $\mathbb{R}$ , not necessarily non-negative, such that  $\int_{\mathbb{R}} K(x)dx = 1$ , and its Fourier transform  $\varphi_K$  is continuously differentiable and supported in  $[-1, 1]$ .*

Both of these assumptions are standard in the literature on deconvolution. Note that Assumption 1 implies that  $f_X$  is bounded and continuous, which in turn implies that  $f_Y$  is bounded and continuous via the relation  $f_Y(y) = \int_{\mathbb{R}} f_X(y-x)f_\varepsilon(x)dx$ .

The next assumption is concerned with the tail behavior of the error characteristic function  $\varphi_\varepsilon$ , which is an important factor that determines the difficulty of estimating  $f_X$  (another factor is the smoothness of  $f_X$ ). We assume here that the error density  $f_\varepsilon$  is ordinary smooth, i.e.,  $|\varphi_\varepsilon(t)|$  decays at most polynomially fast as  $|t| \rightarrow \infty$ .

**Assumption 3.** *The error characteristic function  $\varphi_\varepsilon$  is continuously differentiable and does not vanish on  $\mathbb{R}$ , and there exist constants  $C_1 > 1$  and  $\alpha > 0$  such that  $C_1^{-1}|t|^{-\alpha} \leq |\varphi_\varepsilon(t)| \leq C_1|t|^{-\alpha}$  and  $|\varphi'_\varepsilon(t)| \leq C_1|t|^{-\alpha-1}$  for all  $|t| \geq 1$*

Concrete examples of distributions that satisfy Assumption 3 are Laplace and Gamma distributions together with their convolutions, but apparently many other distributions satisfy Assumption 3. It is not difficult to see that Assumption 3 implies that  $\inf_{|t| \leq h_n^{-1}} |\varphi_\varepsilon(t)| \geq C_1^{-1}(1 - o(1))h_n^\alpha$  as  $n \rightarrow \infty$ .

**Assumption 4.** *Let  $I \subset \mathbb{R}$  be a compact interval such that  $f_Y(y) > 0$  for all  $y \in I$ .*

Now, recall that  $\sigma_n^2(x) = \text{Var}(K_n((x - Y)/h_n))$ . In development of our theory, we will need that  $\sigma_n^2(x)/h_n^{-2\alpha+1}$  is bounded away from zero on  $I$ , on which we would like to make inference on  $f_X$ . It will be shown in Lemma A.3 in the supplementary material that Assumptions 1–4 guarantee that  $\sigma_n^2(x)/h_n^{-2\alpha+1}$  is bounded away from zero on  $I$  for sufficiently large  $n$ .

The next assumption is a mild moment condition on the error distribution, which is used in establishing uniform convergence rates of the empirical characteristic function (see Lemma B.1 in the supplementary material).

**Assumption 5.**  $E[|\varepsilon|^p] < \infty$  for some  $p > 0$ .

The next assumption (mildly) restricts the bandwidth  $h_n$  and the sample size  $m = m_n$  for  $f_\varepsilon$ .

**Assumption 6.** (a)  $n^{-1/2}h_n^{-1} = o\{(\log(1/h_n))^{-1}\}$ . (b)  $m = m_n \rightarrow \infty$  as  $n \rightarrow \infty$ , and

$$m^{-1} = o\left[\{(nh_n \log(1/h_n))^{-1} \wedge \{h_n^{2\alpha+2}(\log(1/h_n))^{-2}\}\right].$$

**Remark 3.1.** For an illustrative purpose, consider the canonical case where  $m = n$ . Then Assumption 6 reduces to the following simple condition:

$$nh_n^{2+2\alpha}/(\log(1/h_n))^2 \rightarrow \infty. \quad (3.1)$$

The conventional “optimal” bandwidth that minimizes the MISE of the kernel estimator (when  $f_\varepsilon$  is known) is proportional to  $n^{-1/(2\alpha+2\beta+1)}$  where  $\beta$  is the “smoothness” of  $f_X$  (cf. Fan, 1991a), and so condition (3.1) is satisfied with this bandwidth if  $\beta > 1/2$ . See also Corollary 3.2 below.

The following theorem establishes that the distribution of the supremum in absolute value of the stochastic process  $\widehat{Z}_n(x)$ ,  $x \in I$  can be approximated by that of a tight Gaussian random variable  $Z_n^G$  in  $\ell^\infty(I)$  with mean zero and the same covariance function as  $Z_n^*$ . This theorem is a building block for establishing validity of the Gaussian multiplier bootstrap described in the previous section. Recall that a Gaussian process  $Z = \{Z(x) : x \in I\}$  indexed by  $I$  is a tight random variable in  $\ell^\infty(I)$  if and only if  $I$  is totally bounded for the intrinsic pseudo-metric  $\rho_2(x, y) = \sqrt{\mathbb{E}[\{Z(x) - Z(y)\}^2]}$  for  $x, y \in I$ , and  $Z$  has sample paths almost surely uniformly  $\rho_2$ -continuous; see van der Vaart and Wellner (1996, p.41).

**Theorem 3.1.** *Under Assumptions 1–6, for each sufficiently large  $n$ , there exists a tight Gaussian random variable  $Z_n^G$  in  $\ell^\infty(I)$  with mean zero and the same covariance function as  $Z_n^*$ , and such that as  $n \rightarrow \infty$ ,  $\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\widehat{Z}_n\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \rightarrow 0$ .*

When  $I$  is not singleton, it is possible to further show that  $\|Z_n^G\|_I$  (and hence  $\|\widehat{Z}_n\|_I$ ) properly normalized converges in distribution to a Gumbel distribution (i.e., a Smirnov-Bickel-Rosenblatt type limit theorem) under additional (substantial) conditions, as in Bisantz et al. (2007). However, we intentionally stop at the “intermediate” Gaussian approximation and do not derive the Gumbel approximation, because of the following two reasons.

1) The Gumbel approximation is poor. The coverage error of the resulting confidence band is of order  $1/\log n$  (Hall, 1991). 2) Deriving the Gumbel approximation requires additional substantial conditions. Because of the slow rate of the Gumbel approximation, it is often preferred to use versions of bootstraps to construct confidence bands for nonparametric density and regression functions (see, e.g., Claeskens and van Keilegom, 2003; Bissantz et al., 2007), but the Gumbel approximation was used as a building block for showing validity of the bootstraps. It was, however, pointed out in Chernozhukov et al. (2014b) that the intermediate Gaussian approximation (such as that in Theorem 3.1) is sufficient for showing validity of bootstraps. We defer the discussion on the regularity conditions to the end of this section.

Another technicality in the proof of Theorem 3.1 is to bound the estimation error of  $\widehat{\varphi}_\varepsilon$ . Dattner et al. (2016, p.172) derive a bound on  $\|\widehat{f}_X - \widehat{f}_X^*\|_{\mathbb{R}}$  that is of order  $O_{\mathbb{P}}\{h_n^{-\alpha}(mh_n)^{-1/2}\}$ , but this rate is not sufficient for our purpose and in particular excludes the case with  $m = n$  in Theorem 3.1; see Step 2 in the proof of Theorem 3.1. Hence to bound the estimation error of  $\widehat{\varphi}_\varepsilon$ , we require a novel idea beyond Dattner et al. (2016).

As a byproduct of the techniques used to prove Theorem 3.1, we can derive uniform convergence rates of  $\widehat{f}_X$  on  $\mathbb{R}$ . In the next corollary, Assumption 4 is not needed.

**Corollary 3.1.** *Suppose that Assumptions 1–3, 5, and 6 are satisfied. Then  $\|\widehat{f}_X(\cdot) - E[\widehat{f}_X^*(\cdot)]\|_{\mathbb{R}} = O_{\mathbb{P}}\{h_n^{-\alpha}(nh_n)^{-1/2}\sqrt{\log(1/h_n)}\}$  as  $n \rightarrow \infty$ .*

Corollary 3.1 does not take into account the bias  $\|E[\widehat{f}_X^*(\cdot)] - f_X(\cdot)\|_{\mathbb{R}}$ , but the above rate is the correct one for the “variance part” when  $f_\varepsilon$  is known. The situation here is bit complicated since  $f_\varepsilon$  is unknown and because of the role of  $m$ . So we only present the following corollary, which deals with the canonical case where  $m = n$ . In the following, for  $\beta > 0$  and  $B > 0$ , let  $\Sigma(\beta, B)$  denote a Hölder ball of functions on  $\mathbb{R}$  with smoothness  $\beta$

and radius  $B$ , namely,

$$\Sigma(\beta, B) = \{f : \mathbb{R} \rightarrow \mathbb{R} : f \text{ is } k\text{-times differentiable and} \\ |f^{(k)}(x) - f^{(k)}(y)| \leq B|x - y|^{\beta-k} \forall x, y \in \mathbb{R}\},$$

where  $k$  is the integer such that  $k < \beta \leq k + 1$ .

**Corollary 3.2.** *Suppose that Assumptions 1–3 and 5 are satisfied, and consider the case where  $m = n$ . Further, suppose that  $f_X \in \Sigma(\beta, B)$  for some  $\beta > 1/2, B > 0$ , and that  $\varphi_K$  is  $(k + 3)$ -times continuously differentiable with  $\varphi_K^{(\ell)}(0) = 0$  for all  $\ell = 1, \dots, k$  where  $k$  is the integer such that  $k < \beta \leq k + 1$ . Take  $h_n = C(n/\log n)^{-1/(2\alpha+2\beta+1)}$  for any constant  $C > 0$ ; then  $\|\widehat{f}_X - f_X\|_{\mathbb{R}} = O_{\mathbb{P}}\{(n/\log n)^{-\beta/(2\alpha+2\beta+1)}\}$ .*

**Remark 3.2.** Informally, for a given error density  $f_{\varepsilon}$  such that  $|\varphi_{\varepsilon}(t)|$  decays like  $|t|^{-\alpha}$  as  $|t| \rightarrow \infty$ ,  $(n/\log n)^{-\beta/(2\alpha+2\beta+1)}$  is the minimax rate of convergence for estimating  $f_X$  under the sup-norm loss when  $f_X \in \Sigma(\beta, B)$  and there is no additional sample from the error distribution. See Theorem 1 in Lounici and Nickl (2011) for the precise formulation. In fact, the proof of Theorem 1 in Lounici and Nickl (2011) continues to hold even when there is a sample  $(\eta_1, \dots, \eta_n)$  from the error distribution that is independent from  $(Y_1, \dots, Y_n)$  – in their proof, modify  $P_k^n$  to be the distribution admitting the joint density  $\prod_{j=1}^n (f_k * f_{\varepsilon})(y_j) f_{\varepsilon}(\eta_j)$ . Hence Corollary 3.2 shows that  $\widehat{f}_X$  attains the minimax rate under the sup-norm loss for  $\beta > 1/2$ , provided that other technical conditions are satisfied.

**Remark 3.3.** The literature on uniform convergence rates in deconvolution is limited. Lounici and Nickl (2011) and Giné and Nickl (2016, Section 5.3.2) derive uniform convergence rates for deconvolution wavelet and kernel density estimators on the entire real line assuming that the error density is known; Dattner et al. (2016, Proposition 2.6) derive uniform convergence rates for the deconvolution kernel density estimator with the estimated

error characteristic function, but on a bounded interval. So their results do not cover the above corollaries, although combination of a bound on  $\|\widehat{f}_X - \widehat{f}_X^*\|_{\mathbb{R}}$  in the proof of Proposition 2.6 in Dattner et al. (2016) with the results in Giné and Nickl (2016, Section 5.3.2) would yield Corollary 3.2.

Now, we present validity of the multiplier bootstrap confidence band described in the previous section.

**Theorem 3.2.** *Under Assumptions 1–6, as  $n \rightarrow \infty$ ,*

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\widehat{Z}_n^\xi\|_I \leq z \mid \mathcal{D}_n\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \xrightarrow{\mathbb{P}} 0, \quad (3.2)$$

where  $\mathcal{D}_n = \{Y_1, \dots, Y_n, \eta_1, \dots, \eta_m\}$  and  $Z_n^G$  is the Gaussian random variable in  $\ell^\infty(I)$  given in Theorem 3.1. Therefore, if we denote by  $\widehat{c}_n(1 - \tau)$  the  $(1 - \tau)$ -quantile of the conditional distribution of  $\|\widehat{Z}_n^\xi\|_I$  given  $\mathcal{D}_n$ , then we have

$$\mathbb{P}\{\|\widehat{Z}_n^\xi\|_I \leq \widehat{c}_n(1 - \tau)\} \rightarrow 1 - \tau \quad (3.3)$$

as  $n \rightarrow \infty$ . Finally, if we denote by  $s_n(x)$  the width of the band  $\widehat{\mathcal{C}}_n(x)$  for each  $x \in I$ , then  $\sup_{x \in I} s_n(x) = O_{\mathbb{P}}\{h_n^{-\alpha}(nh_n)^{-1/2}\sqrt{\log(1/h_n)}\}$ .

Theorem 3.2 shows that the multiplier bootstrap confidence band  $\widehat{\mathcal{C}}_n$  defined in (2.5) contains the surrogate function  $\mathbb{E}[\widehat{f}_X^*(\cdot)]$  on  $I$  with probability  $1 - \tau + o(1)$  as  $n \rightarrow \infty$ . If  $f_X$  belongs to a Hölder ball  $\Sigma(\beta, B)$ , then  $\widehat{\mathcal{C}}_n$  will be a valid confidence band for  $f_X$  provided that  $h_n$  is chosen in such a way that  $h_n^{\alpha+\beta}\sqrt{nh_n \log(1/h_n)} \rightarrow 0$ , which corresponds to choosing undersmoothing bandwidths.

**Corollary 3.3.** *Suppose that Assumptions 1–6 are satisfied. Further, suppose that  $f_X \in \Sigma(\beta, B)$  for some  $\beta > 0, B > 0$ , and that  $\varphi_K$  is  $(k + 3)$ -times continuously differentiable*

with  $\varphi^{(\ell)}(0) = 0$  for all  $\ell = 1, \dots, k$  where  $k$  is the integer such that  $k < \beta \leq k + 1$ . Consider the multiplier bootstrap confidence band  $\widehat{\mathcal{C}}_n$  defined in (2.5). Then as  $n \rightarrow \infty$ ,  $\mathbb{P}\{f_X(x) \in \widehat{\mathcal{C}}_n(x) \forall x \in I\} \rightarrow 1 - \tau$ , provided that

$$h_n^{\alpha+\beta} \sqrt{nh_n \log(1/h_n)} \rightarrow 0. \quad (3.4)$$

Consider the canonical case where  $m = n$ . Then the conditions on the bandwidth  $h_n$  in Corollary 3.3 reduce to  $nh_n^{2\alpha+2}/(\log(1/h_n))^2 \rightarrow \infty$  and  $h_n^{\alpha+\beta} \sqrt{nh_n \log(1/h_n)} \rightarrow 0$ , and so we need  $\beta > 1/2$  in order to ensure the existence of the bandwidth  $h_n$  satisfying these conditions. For example, if  $\beta > 1/2$ , choosing  $h_n = v_n(n/\log n)^{-1/(2\alpha+2\beta+1)}$  for some sequence  $v_n \rightarrow 0$  sufficiently slowly satisfies the above restriction and yields that the supremum width of the band  $\widehat{\mathcal{C}}_n$  is  $\sup_{x \in I} s_n(x) = O_{\mathbb{P}}\{v_n^{-\alpha-1/2}(n/\log n)^{-\beta/(2\alpha+2\beta+1)}\}$ , which can be arbitrarily close to the minimax rate of estimation, namely,  $(n/\log n)^{-\beta/(2\alpha+2\beta+1)}$ .

**Remark 3.4** (Comparisons with Bissantz et al. (2007) and Schmidt-Hieber et al. (2013)). Bissantz et al. (2007) is an important pioneering work on confidence bands in deconvolution. They assume that the error density is known and ordinary smooth, and show that

$$\sqrt{2 \log(1/h_n)} (\|\sqrt{nh_n^{\alpha+1/2}}(\widehat{f}_X^*(\cdot) - \mathbb{E}[\widehat{f}_X^*(\cdot)])/\sqrt{f_Y(\cdot)}\|_{[0,1]}/C_{K,1}^{1/2} - d_n)$$

converges in distribution to a Gumbel distribution, where  $d_n = \sqrt{2 \log(1/h_n)} + \frac{\log(C_{K,2}^{1/2}/2\pi)}{\sqrt{2 \log(1/h_n)}}$ , and where  $C_{K,1}$  and  $C_{K,2}$  are some constants; see Bissantz et al. (2007) for their explicit values. Further, they show validity of the nonparametric bootstrap for approximating the distribution of  $\|\sqrt{nh_n^{\alpha+1/2}}(\widehat{f}_X^*(\cdot) - \mathbb{E}[\widehat{f}_X^*(\cdot)])/\sqrt{f_Y(\cdot)}\|_{[0,1]}$ ; see their Theorem 2.

Since we work with a different setting than Bissantz et al. (2007), namely we assume that  $f_\varepsilon$  is unknown and an auxiliary sample from  $f_\varepsilon$  is available, the regularity conditions in the present paper are not directly comparable to those of Bissantz et al. (2007). However, it is worthwhile to point out that conditions on the error characteristic function are significantly

relaxed in the present paper. Indeed, their Assumption 2 is substantially more restrictive than our Assumption 3. The reasons that they require their Assumption 2 are: 1) they use the KMT strong approximation (Komlós et al., 1975) to the empirical process  $[0, 1] \ni x \mapsto \sqrt{nh_n^{\alpha+1/2}}(\widehat{f}_X^*(x) - \mathbb{E}[\widehat{f}_X^*(x)])/\sqrt{f_Y(x)}$ , to which end a bound on the total variation of  $K_n$  is needed, and their Assumption 2 (a) serves that role; 2) their analysis relies on the Gumbel approximation, to which end, they require further approximations based on extreme value theory (cf Leadbetter et al., 1983) beyond the KMT approximation, and consequently require some extra assumptions, namely, their Assumption 2 (b).

In the present paper, we build upon the intermediate Gaussian and multiplier bootstrap approximation theorems developed in Chernozhukov et al. (2014a,b, 2016), and regularity conditions needed to apply those techniques are typically much weaker than those for the KMT and Gumbel approximations. In particular, we do not need a bound on the total variation of  $K_n$ ; instead, we need that the class of functions  $\{y \mapsto K_n((x-y)/h_n) : x \in I\}$  is of Vapnik-Chervonenkis type, and to that end, thanks to Lemma 1 in Giné and Nickl (2009), it is enough to prove that  $K_n$  has bounded quadratic variation of order  $h_n^{-2\alpha}$ , which is ensured by our Assumptions 2 and 3 (see Lemmas A.1 and A.2 ahead). In addition, in contrast to Bissantz et al. (2007), we do not need that  $\sigma_n^2(x)/h_n^{-2\alpha+1}$  has a fixed limit; we only need that  $\sigma_n^2(x)/h_n^{-2\alpha+1}$  is bounded away from zero uniformly in  $x \in I$ .

Furthermore, the intermediate Gaussian and multiplier bootstrap approximations apply not only to the ordinary smooth case, but also to the super-smooth case, as discussed in Section 6, and so they enable us to study confidence bands for  $f_X$  in a unified way (although in the super-smooth case we require  $m/n \rightarrow \infty$ ). On the other hand, as shown in van Es and Gugushvili (2008), the Gumbel approximation does not hold for the super-smooth case in general (see also Remark 6.1 ahead).

Schmidt-Hieber et al. (2013), assuming that the error density is known and ordinary

smooth, develop methods to make inference on shape constraints for  $f_X$ , which also cover construction of confidence bands (although their main interest is not in confidence bands). They use an intermediate Gaussian approximation different from ours based on the KMT approximation, and are able to relax assumptions in Bissantz et al. (2007) (a minor difference is that they require twice differentiability of  $\varphi_\varepsilon$  while we only require one differentiability). However, the crucial point of their approach is that the distribution of the approximating Gaussian process is known, which is the case when the distribution of  $\varepsilon$  is known but not the case otherwise. Hence their methodology is not directly applicable to our case. We also note that in their methodology,  $f_Y$  appears as a scaling constant, and so that we need to estimate  $f_Y$  separately and thus to choose an appropriate bandwidth for  $f_Y$  separately. On the other hand, we are using a different scaling and separate estimation of  $f_Y$  is not needed.

## 4 Simulation studies

### 4.1 Simulation framework

In this section, we present simulation studies to evaluate finite-sample performance of the inference method developed in the previous two sections. We generate data from the model introduced in Example 2.1. For distributions of the primitive latent variables  $(X, \varepsilon^{(1)}, \varepsilon^{(2)})$ , we consider two alternative models described below.

In the first model,  $X$  is drawn from the centered normal distribution  $N(0, \sigma_X^2)$ , and  $\varepsilon^{(1)}$  and  $\varepsilon^{(2)}$  are drawn from the Laplace distribution with  $(0, 1)$  as the location and scale parameters. This Laplace distribution is symmetric around zero, and therefore the premise of Example 2.1 regarding the error variables is satisfied. The distribution of  $\varepsilon = (\varepsilon^{(1)} +$

	Model 1	Model 2
Distribution of $X$	$N(0, \sigma_X^2)$	$\chi^2(df)$
Distribution of $(\varepsilon^{(1)}, \varepsilon^{(2)})$	Laplace $(0, 1)$	Laplace $(0, \sqrt{2})$
Smoothness of $X$	Super	Ordinary
Smoothness of $\varepsilon$	Ordinary	Ordinary
Signal-to-noise ratio	$\sigma_X$	$\sqrt{df}$
Interval $I$	$[-2\sigma_X, 2\sigma_X]$	$[\mu_X/2, \mu_X + 2\sigma_X]$

Table 1: A summary of the two models considered for simulation studies.

$\varepsilon^{(2)}/2$  has its characteristic function not vanishing on  $\mathbb{R}$  and is ordinary smooth with  $\alpha = 4$ . This setting conveniently yields the signal-to-noise ratio given by

$$\sqrt{\frac{\text{Var}(X)}{\text{Var}(\varepsilon)}} = \sqrt{\frac{\sigma_X^2}{\text{Var}(\varepsilon^{(1)})/4 + \text{Var}(\varepsilon^{(2)})/4}} = \sigma_X.$$

In the second model,  $X$  is drawn from the chi-squared distribution  $\chi^2(df)$ , and  $\varepsilon^{(1)}$  and  $\varepsilon^{(2)}$  are drawn from the Laplace distribution with  $(0, \sqrt{2})$  as the location and scale parameters. The distribution of  $\varepsilon = (\varepsilon^{(1)} + \varepsilon^{(2)})/2$  has its characteristic function not vanishing on  $\mathbb{R}$  and is ordinary smooth with  $\alpha = 4$ . In this setting, the signal-to-noise ratio is given by

$$\sqrt{\frac{\text{Var}(X)}{\text{Var}(\varepsilon)}} = \sqrt{\frac{2df}{\text{Var}(\varepsilon^{(1)})/4 + \text{Var}(\varepsilon^{(2)})/4}} = \sqrt{df}.$$

Table 1 summarizes the two models and their relevant properties.

The observed portion of data,  $\mathcal{D}_n = \{Y_1, \dots, Y_n, \eta_1, \dots, \eta_n\}$ , is constructed by  $Y_j = (Y_j^{(1)} + Y_j^{(2)})/2$  and  $\eta_j = (Y_j^{(1)} - Y_j^{(2)})/2$ , where  $Y_j^{(1)} = X_j + \varepsilon_j^{(1)}$  and  $Y_j^{(2)} = X_j + \varepsilon_j^{(2)}$ , for each  $j = 1, \dots, n$ . The three primitive latent variables,  $X$ ,  $\varepsilon^{(1)}$ , and  $\varepsilon^{(2)}$  are independently generated. We use Monte Carlo simulations to compute the coverage probabilities of our

multiplier bootstrap confidence bands for  $f_X$  on the interval  $I = [-2\sigma_X, 2\sigma_X]$  for Model 1 and on the interval  $I = [\mu_X/2, \mu_X + 2\sigma_X]$  where  $(\mu_X, \sigma_X) = (df, \sqrt{2df})$  for Model 2. We use the kernel function  $K$  defined by its Fourier transform  $\varphi_K$  as follows:

$$\varphi_K(t) = \begin{cases} 1 & \text{if } |t| \leq c \\ \exp\left\{\frac{-b \exp(-b/(|t|-c)^2)}{(|t|-1)^2}\right\} & \text{if } c < |t| < 1, \\ 0 & \text{if } |t| \geq 1 \end{cases}$$

where  $b = 1$  and  $c = 0.05$  (cf. Politis and Romano, 1999; Bissantz et al., 2007). Note that  $\varphi_K$  is supported in  $[-1, 1]$ , and its inverse Fourier transform  $K$  is real-valued and integrable with  $\int_{\mathbb{R}} K(x)dx = 1$ . For the bandwidth selection, we follow a data-driven rule discussed in the next subsection, inspired by Bissantz et al. (2007).

## 4.2 Bandwidth selection

Our theory prescribes admissible asymptotic rates for the bandwidth  $h_n$  that require under-smoothing. The literature provides data-driven approaches to bandwidth selection, which are usually based on minimizing the MISE. These data-driven approaches tend to yield non-under-smoothing bandwidths, and do not conform with our requirements. We adopt the two-step selection method developed in Bissantz et al. (2007, Section 5.2) that aims to select undersmoothing bandwidths. The first step selects a pilot bandwidth  $h_n^P$  based on a data-driven approach. We simply use a normal reference bandwidth (Delaigle and Gijbels, 2004, Section 3.1) for  $h_n^P$ . Once the pilot bandwidth  $h_n^P$  is obtained, we next make a list of candidate bandwidths  $h_{n,j} = (j/J)h_n^P$  for  $j = 1, \dots, J$ . The deconvolution estimate based on the  $j$ -th candidate bandwidth is denoted by  $\widehat{f}_{X,j}$ . The second step in the two step approach chooses the largest bandwidth  $h_{n,j}$  such that the adjacent uniform distance  $\|\widehat{f}_{X,j-1} - \widehat{f}_{X,j}\|_I$  is larger than  $\rho\|\widehat{f}_{X,J-1} - \widehat{f}_{X,J}\|_I$  in the pilot case for some  $\rho > 1$ . Similarly

Nominal Coverage Probability ( $1 - \tau$ )	Sample Size ( $n$ )	(A) Model 1			(B) Model 2		
		Signal-to-Noise Ratio			Signal-to-Noise Ratio		
		2.0	4.0	8.0	2.0	4.0	8.0
0.800	250	0.762	0.755	0.728	0.636	0.740	0.666
	500	0.786	0.746	0.763	0.698	0.751	0.685
	1,000	0.784	0.750	0.754	0.732	0.749	0.702
0.900	250	0.870	0.866	0.843	0.760	0.853	0.799
	500	0.897	0.862	0.863	0.822	0.867	0.822
	1,000	0.897	0.862	0.863	0.848	0.878	0.836
0.950	250	0.930	0.936	0.907	0.834	0.927	0.871
	500	0.951	0.929	0.923	0.834	0.928	0.897
	1,000	0.942	0.933	0.933	0.912	0.940	0.906

Table 2: Simulated uniform coverage probabilities of  $f_X$  by estimated confidence bands in  $I = [-2\sigma_X, 2\sigma_X]$  for Model 1 and  $I = [\mu_X/2, \mu_X + 2\sigma_X]$  for Model 2. The simulated probabilities are computed for each of the three nominal coverage probabilities, 80%, 90%, and 95%, based on 2,000 Monte Carlo iterations.

to the values recommended by Bissantz et al. (2007), we find that  $J \approx 20$  and  $\rho \approx 3$  work well in our simulation studies.

### 4.3 Simulation results

Simulated uniform coverage probabilities are computed for each of the three nominal coverage probabilities, 80%, 90%, and 95%, based on 2,000 Monte Carlo iterations. In each run of the simulation, we generate 2,500 multiplier bootstrap replications given the observed data  $\mathcal{D}_n$  to compute the estimated critical values,  $\hat{c}_n(1 - \tau)$ .

Results under Models 1 and 2 are summarized in column groups (A) and (B), respectively, of Table 2 for each of the three different cases of the signal-to-noise ratio:  $\sigma_X \in \{2.0, 4.0, 8.0\}$ , and for each of the three sample sizes  $n = m \in \{250, 500, 1,000\}$ . Observe that the simulated probabilities are close to the respective nominal probabilities. Not surprisingly, the size tends to be more accurate for the results based on larger sample sizes. The simulated probabilities are closer to the nominal probabilities in (A) than (B).


## 5 Application to OCS Wildcat auctions

In this section, we apply our method to the Outer Continental Shelf (OCS) Auction Data (see Hendricks et al. (1987) for details), and construct a confidence band for the density of mineral rights on oil and gas on offshore lands off the coasts of Texas and Louisiana in the gulf of Mexico. We focus on “wildcat sales,” referring to sales of those oil and gas tracts whose geological or seismic characteristics are unknown to participating firms. The sales rule follows the first-price sealed-bid auction mechanism, where participating firms simultaneously submit sealed bids, and the highest bidder pays the price they submitted to receive the right for the tract. Firms who are willing to participate in sales can carry out a seismic investigation before the sales date in order to estimate the value of mineral rights. The ex ante value  $Y^{(1)}$  (in the logarithm of US dollars per acre) obtained by firm 1 through its investigation of the tract is treated as a measure of the ex post value  $X$  (also known as the common component, in the logarithm of US dollars per acre) with an assessment error  $\varepsilon^{(1)}$  (also known as the private component), i.e.,  $Y^{(1)} = X + \varepsilon^{(1)}$ . Collecting the ex ante values  $(Y^{(1)}, Y^{(2)})$  for pairs of firms across various wildcat auctions, we can obtain data necessary to construct a uniform confidence band for the density  $f_X$  of ex post mineral right values under our assumptions.

For this setup and for this data set, Li et al. (2000) apply the method of Li and Vuong (1998) to nonparametrically estimate  $f_X$ , but they do not obtain a confidence band. In their analysis, firms' ex ante values  $(Y^{(1)}, Y^{(2)})$  are first recovered from bid data through a widely used method in economics which is based on an equilibrium restriction (Bayesian Nash equilibrium) for the first-price sealed-bid auction mechanism – see our supplementary material. In this paper, we directly take these ex ante values  $(Y^{(1)}, Y^{(2)})$  as the data to be used as an input for our analysis. The sample consists of 169 tracts with 2 firms in each tract. We next construct our auxiliary data  $(Y, \eta)$  following Example 2.1.

We continue to use the same kernel function and the same bandwidth selection rule as those ones used for simulation studies in Section 4. Confidence bands for  $f_X$  are constructed using 25,000 multiplier bootstrap replications. Figure 1 (A) shows the obtained 90% and 95% confidence bands in light gray and dark gray, respectively. The black curve draws our nonparametric estimate of  $f_X$ , and runs at the center of the bands. Our nonparametric estimate, our confidence bands, and the nonparametric estimate obtained by Li et al. (2000) – shown in their Figure 4 – are all similar to each other, and share the same qualitative characteristics. First, unlike the bimodal density of the private values  $(Y^{(1)}, Y^{(2)})$  (see Figure 2 in the supplementary material), the density of the common value  $X$  is suggested to be single-peaked in all the results of our analysis and that of Li et al. (2000). Second, as Li et al. (2000) also emphasized, the small bump in the estimated density  $f_X$  around  $x = 6$  is common in all the results. Furthermore, our confidence bands do not include zero at this locality,  $x = 6$ . This result can be viewed as a statistical evidence in support of a significant presence of such a bump pointed out by Li et al. (2000). A more global look into the graph suggests that the 95% confidence band is bounded away from zero on the interval  $[3.90, 6.32]$ . In other words, we can conclude that the ex post value of mineral rights in the US dollars per acre is supported on a superset of the interval  $[\exp(3.90), \exp(6.32)] \approx$

(A) Unknown Error Distribution



(B) Laplace Error Distribution




Figure 1: 90% and 95% confidence bands for  $f_X$  marked in light gray and dark gray, respectively. The black curve in the middle indicates our deconvolution kernel density estimate of  $f_X$ .

[49, 555], if we consider a set of density functions contained in the 95% confidence band.

Finally, we show an empirical result that we would obtain under the assumption that the distribution of  $\varepsilon$  were known as in the previous literature. Specifically, for this exercise, we assume that  $\varepsilon$  follows  $\text{Laplace}(0, \sqrt{\sigma_\varepsilon^2/2})$ , where  $\sigma_\varepsilon^2 = 0.168$  – this number is the sample variance of  $\eta_j$ . Figure 1 (B) shows the result. There are some notable differences from Figure 1 (A). Most importantly, the 95% confidence band now contains zero around the locality,  $x = 6$ , of the aforementioned bump in the estimated density of  $X$ . The contrast between Figure 1 (A) and (B) shows that there can be non-trivial differences in statistical implications of confidence bands between the case of assuming known error distribution and the case of assuming unknown error distribution.

## 6 Extensions to super-smooth case

In this section, we consider extensions of the results on confidence bands to the case where the error density  $f_\varepsilon$  is super-smooth. We still keep Assumptions 1 and 5, but require a different set of assumptions on the kernel  $K$ , the bandwidth  $h_n$ , and the sample size  $m$  for  $f_\varepsilon$ . It turns out that from a technical reason, we require  $m/n \rightarrow \infty$  in the super-smooth case, and so Example 2.1 is formally not covered in the super-smooth case. Still, we believe that the extensions to the super-smooth case are of some interest.

We modify Assumptions 2, 3, and 6 as follows. First, for the kernel and error characteristic functions, we assume the following conditions.

**Assumption 7.** *Let  $K$  be a kernel on  $\mathbb{R}$  such that its Fourier transform  $\varphi_K$  is even (i.e.,  $\varphi_K(-t) = \varphi_K(t)$ ) and supported in  $[-1, 1]$ . Furthermore, there exist constants  $C_2 \neq 0$  and  $\lambda \geq 0$  such that  $\varphi_K(1-t) = C_2 t^\lambda + o(t^\lambda)$  as  $t \downarrow 0$ .*

**Assumption 8.** *The error characteristic function  $\varphi_\varepsilon$  does not vanish on  $\mathbb{R}$ , and there exist constants  $C_3 \in \mathbb{R} \setminus \{0\}$ ,  $\gamma > 1$ ,  $\gamma_0 \in \mathbb{R}$ ,  $\nu > 0$  such that  $\varphi_\varepsilon(t) = C_3(1 + o(1))|t|^{\gamma_0} e^{-\nu|t|^\gamma}$  as  $|t| \rightarrow \infty$ .*

These assumptions are adapted from van Es and Uh (2005). Assumption 8 covers cases where the error characteristic function decays exponentially fast as  $|t| \rightarrow \infty$ , thereby covering cases where the error density is super-smooth. However, Assumption 8 is more restrictive than standard super-smoothness conditions; e.g., it excludes the Cauchy error. This assumption is needed to derive a lower bound on  $\sigma_n^2(x)$ ; see the following discussion.

Assumptions 7 and 8, together with the assumption that the variance function  $E[Y^2] < \infty$ , ensure that  $\sigma_n^2(x) = \text{Var}(K_n(x - Y)/h_n)$  is expanded as

$$\sigma_n^2(x) = (1 + o(1)) \frac{C_2^2}{2C_3^2\pi^2} (\nu\gamma)^{-2\lambda-2} (\Gamma(\lambda+1))^2 h_n^{2\gamma(1+\lambda)+2\gamma_0} e^{2\nu h_n^{-\gamma}} \quad (6.1)$$

as  $n \rightarrow \infty$ ; see the proof of Theorem 1.5 in van Es and Uh (2005). It is not difficult to verify from their proof that  $o(1)$  in (6.1) is uniform in  $x \in I$  for any compact interval  $I \subset \mathbb{R}$ . It is worthwhile to point out that, in contrast to the ordinary smooth case, the lower bound on  $\sigma_n^2(x)$  in (6.1) does not explicitly depend on  $x$  nor  $f_Y$ . Further, Assumption 8 implies that

$$\inf_{|t| \leq h_n^{-1}} |\varphi_\varepsilon(t)| \geq |C_3|(1 - o(1))h_n^{-\gamma_0} e^{-\nu h_n^{-\gamma}} \quad (6.2)$$

as  $n \rightarrow \infty$ . It turns out that (6.1) and (6.2) are the only differences to take care of when proving the analogues of Theorems 3.1 and 3.2 in the super-smooth case. Finally, we modify Assumption 6 as follows.

**Assumption 9.** (a)  $n^{-1/2}h_n^{-2\gamma(1+\lambda)} = o\{(\log(1/h_n))^{-1}\}$ . (b)  $m = m_n \rightarrow \infty$  as  $n \rightarrow \infty$ , and

$$m^{-1} = o\left[\left\{n^{-1}h_n^{2\gamma(1+\lambda)-2}(\log(1/h_n))^{-1}\right\} \wedge \left\{h_n^{4\gamma(1+\lambda)-2\gamma_0}e^{-2\nu h_n^{-\gamma}}(\log(1/h_n))^{-2}\right\}\right].$$

The requirement that  $m^{-1} = o\{n^{-1}h_n^{2\gamma(1+\lambda)-2}(\log(1/h_n))^{-1}\}$  implies that we at least need  $m/n \rightarrow \infty$ . This condition is used to ensure that the effect of estimating  $\varphi_\varepsilon$  is negligible. To be precise, in our proof, a bound on  $\|\widehat{f}_X - \widehat{f}_X^*\|_{\mathbb{R}}$  involves a term of order  $m^{-1/2}h_n^{\gamma_0}e^{\nu h_n^{-\gamma}}$ , which has to be of smaller order than  $n^{-1/2}h_n^{\gamma(1+\lambda)+\gamma_0-1}e^{\nu h_n^{-\gamma}}(\log(1/h_n))^{-1/2}$ . Technically, this problem happens because the ratio of  $1/\inf_{|t| \leq h_n^{-1}} |\varphi_\varepsilon(t)|$  over  $\inf_{x \in I} \sigma_n(x)$  is larger in the super-smooth case than that in the ordinary smooth case; the ratio is  $O(h_n^{-\gamma(1+\lambda)})$  in the super-smooth case, while it is  $O(h_n^{-1/2})$  in the ordinary smooth case. It is not known at the current moment whether we could relax this condition on  $m$  in the super-smooth case.

In any case, these assumptions guarantee that the conclusions of Theorems 3.1 and 3.2, except for the result on the width of the band, hold true in the super-smooth case.

**Theorem 6.1.** *Suppose that Assumptions 1, 5, and 7–9 are satisfied. Let  $I \subset \mathbb{R}$  be any compact interval, and suppose in addition that  $\mathbb{E}[Y^2] < \infty$ . Then the conclusions of Theorems 3.1 and 3.2, except for the result on the width of the band, hold true.*

**Remark 6.1** (Comparisons with van Es and Gugushvili (2008)). van Es and Gugushvili (2008) prove that, under the assumptions that  $f_\varepsilon$  is known and satisfies Assumption 8 (of the present paper) with  $\gamma = 2$ ,

$$\frac{\sqrt{n}}{h_n^{2(1+\lambda)+\gamma_0-1} e^{\nu h_n^{-2}}} \|\widehat{f}_X^*(\cdot) - \mathbb{E}[\widehat{f}_X^*(\cdot)]\|_{[0,1]} \xrightarrow{d} \frac{|C_2|}{\sqrt{2}|C_3|\pi} (2\nu)^{-\lambda-1} \Gamma(\lambda+1)V,$$

where  $V$  follows the Rayleigh distribution, i.e.,  $V$  is a random variable having density  $f_V(v) = ve^{-v^2/2} 1_{[0,\infty)}(v)$  (see van Es and Gugushvili, 2008, for the precise regularity conditions). Interestingly, the limit distribution differs from Gumbel distributions.

Despite this non-standard feature, Theorem 6.1 shows that the multiplier bootstrap “works”, i.e., the conditional distribution of  $\|\widehat{Z}_n^\xi\|_I$  can consistently estimate the distribution of  $\|\widehat{Z}_n\|_I$  in the sense that  $\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\widehat{Z}_n^\xi\|_I \leq z \mid \mathcal{D}_n\} - \mathbb{P}\{\|\widehat{Z}_n\|_I \leq z\}| \xrightarrow{P} 0$ . Further, Theorem 6.1 extends the admissible range of  $\gamma$  compared with the result of van Es and Gugushvili (2008).

If  $f_X$  belongs to a Hölder ball  $\Sigma(\beta, B)$ , then we have the following corollary.

**Corollary 6.1.** *Assume all the conditions of Theorem 6.1. Further, suppose that  $f_X \in \Sigma(\beta, B)$  for some  $\beta > 0, B > 0$ , and that  $\varphi_K$  is  $(k+3)$ -times continuously differentiable with  $\varphi_K^{(\ell)}(0) = 0$  for all  $\ell = 1, \dots, k$  where  $k$  is the integer such that  $\beta < k \leq \beta + 1$ . Consider the multiplier bootstrap confidence band  $\widehat{\mathcal{C}}_n$  defined in (2.5). Then as  $n \rightarrow \infty$ ,  $\mathbb{P}\{f_X(x) \in \widehat{\mathcal{C}}_n(x) \forall x \in I\} \rightarrow 1 - \tau$ , provided that  $h_n^{\beta+1-\gamma(1+\lambda)-\gamma_0} e^{-\nu h_n^{-\gamma}} \sqrt{n \log(1/h_n)} \rightarrow 0$ .*

## SUPPLEMENTARY MATERIAL

**Supplementary material:** The supplementary material contains all the proofs, additional simulation results, and a description on the empirical data.

## References

- Bickel, P. and Rosenblatt, M. (1973). On some global measures of the deviations of density function estimates. *Ann. Statist.* **1** 1071-1095. Correction (1975) **3** 1370.
- Bissantz, N., Dümbgen, L., Holzmann, H., and Munk, A. (2007). Non-parametric confidence bands in deconvolution density estimation. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* **69** 483-506.
- Bissantz, N. and Holzmann, H. (2008). Statistical inference for inverse problems. *Inverse Problems* **24**:034009.
- Bohnomme, S. and Robin, J.-M. (2010). Generalized nonparametric deconvolution with an application to earnings dynamics. *Rev. Econom. Stud.* **77** 491-533.
- Carroll, R.J. and Hall, P. (1988). Optimal rates of convergence for deconvolving a density. *J. Amer. Statist. Assoc.* **83** 1184-1186.
- Carroll, R.J., Ruppert, D., Stefanski, L.A., and Crainiceanu, C.M. (2006). *Measurement Error in Nonlinear Models: A Modern Perspective (2nd Edition)*. Chapman & Hall/CRC.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014a). Gaussian approximation of suprema of empirical processes. *Ann. Statist.* **42** 1564-1597.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014b). Anti-concentration and honest, adaptive confidence bands. *Ann. Statist.* **42** 1787-1818.

- Chernozhukov, V., Chetverikov, D., and Kato, K. (2016). Empirical and multiplier bootstraps for suprema of empirical processes of increasing complexity, and related Gaussian couplings. *Stochastic Process. Appl.*, to appear. arXiv:1502:00352.
- Claeskens, G. and van Keilegom, I. (2003). Bootstrap confidence bands for regression curves and their derivatives. *Ann. Statist.* **31** 1852-1884.
- Comte, F. and Lacour, C. (2011). Data-driven density estimation in the presence of additive noise with unknown distribution. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* **73** 601-627.
- Dattner, I., Reiß, M., and Trabs, M. (2016). Adaptive quantile estimation in deconvolution with unknown error distribution. *Bernoulli* **22** 143192.
- Delaigle, A. and Gijbels, I. (2004). Practical bandwidth selection in deconvolution kernel density estimation. *Comput. Statist. Data Anal.* **45** 249-267.
- Delaigle, A. and Hall, P. (2016). Methodology for nonparametric deconvolution when the error distribution is unknown. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* **78** 231-252.
- Delaigle, A., Hall, P., and Jamshidi, F. (2015). Confidence bands in nonparametric errors-in-variables regression. *J. R. Stat. Soc. Ser. B. Stat. Methodol.* **77** 149-169.
- Delaigle, A., Hall, P., and Meister, A. (2008). On deconvolution with repeated measurements. *Ann. Statist.* **36** 665-685.
- Diggle, P.J. and Hall, P. (1993). A Fourier approach to nonparametric deconvolution of a density estimate. *J. Roy. Stat. Soc. Ser. B. Stat. Methodol.* **55** 523-531.
- Efromovich, S. (1997). Density estimation for the case of supersmooth measurement error. *J. Amer. Stat. Assoc.* **92** 526-535.

- van Es, B. and Gugushvili, S. (2008). Weak convergence of the supremum distance for supersmooth kernel deconvolution. *Statist. Probab. Lett.* **78** 2932-2938.
- van Es, B. and Uh, H.-W. (2005). Asymptotic normality of kernel-type deconvolution estimators. *Scand. J. Statist.* **32** 467-483.
- Fan, J. (1991a). On the optimal rates of convergence for nonparametric deconvolution problems. *Ann. Statist.* **19** 1257-1272.
- Fan, J. (1991b). Asymptotic normality for deconvolution kernel density estimators. *Sankhya A* **53** 97-110.
- Folland, G.B. (1999). *Real Analysis (2nd Edition)*. Wiley.
- Giné, E. and Nickl, R. (2009). Uniform limit theorems for wavelet density estimators. *Ann. Probab.* **37** 1605-1646.
- Giné, E. and Nickl, R. (2016). *Mathematical Foundations of Infinite-Dimensional Statistical Models*. Cambridge University Press.
- Hall, P. (1991). On convergence rates of suprema. *Probab. Theory Related Fields* **89** 447-455.
- Hendricks, K., Porter, R.H., and Boudreau, B. (1987). Information, returns, and bidding behavior in OCS auctions: 1954-1969. *J. Indust. Econom.* **35** 517-542.
- Horowitz, J.L. (2009). *Semiparametric and Nonparametric Methods in Econometrics*. Springer.
- Horowitz, J.L. and Markatou, M. (1996). Semiparametric estimation of regression models for panel data. *Rev. Econom. Stud.* **63** 145-168.
- Johannes, J. (2009). Deconvolution with unknown error distribution. *Ann. Statist.* **37** 2301-2323.

- Komlós, J., Major, P., and Tusnády, G. (1975). An approximation for partial sums of independent rv's and the sample df I. *Z. Warhsch. Verw. Gabiete* **32** 111-131.
- Krasnokutskaya, E. (2011). Identification and estimation of auction models with unobserved heterogeneity. *Rev. Econ. Stud.* **78** 293-327.
- Leadbetter, M.R., Lindgren, G., and Rootzén, H. (1983). *Extremes and Related Properties of Random Sequences and Processes*. Springer.
- Li, T., Perrigne, I., and Vuong, Q. (2000). Conditionally independent private information in OCS wildcat auctions. *J. Econometrics* **98** 129-161.
- Li, T. and Vuong, Q. (1998). Nonparametric estimation of the measurement error model using multiple indicators. *J. Multivariate Anal.* **65** 139-165.
- Lounici, K. and Nickl, R. (2011). Global uniform risk bounds for wavelet deconvolution estimators. *Ann. Statist.* **39** 201-231.
- Meister, A. (2009). *Deconvolution Problems in Nonparametric Statistics*. Springer.
- Neumann, M.H. (1997). On the effect of estimating the error density in nonparametric deconvolution. *J. Nonparametric Statist.* **7** 307-330.
- Neumann, M.H. (2007). Deconvolution from panel data with unknown error distribution. *J. Multivariate Anal.* **98** 1955-1968.
- Neumann, M.H. and Reiß, M. (2009). Nonparametric estimation for Lévy processes from low-frequency observations. *Bernoulli* **15** 223-248.
- Politis, D.N. and Romano, J.P. (1999). Multivariate density estimation with general flat-top kernels of infinite order. *J. Multivariate Anal.* **68** 1-25.
- Schennach, S.M. (2013). Convolution without independence. Cemmap working paper.

- Schmidt-Hieber, J., Munk, A., and Dümbgen, L. (2013). Multiscale methods for shape constraints in deconvolution: confidence statements for qualitative features. *Ann. Statist.* **41** 1299-1328.
- Smirnov, N.V. (1950). On the construction of confidence regions for the density of distribution of random variables. *Doklady Akad. Nauk SSSR* **74** 189-191 (Russian).
- Stefanski, L. and Carroll, R.J. (1990). Deconvoluting kernel density estimators. *Statistics* **21** 169-184.
- van der Vaart, A.W. and Wellner, J.A. (1996). *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer.

# Supplementary material for “Uniform confidence bands in deconvolution with unknown error distribution”

Kengo Kato\*

Graduate School of Economics, University of Tokyo

and

Yuya Sasaki

Department of Economics, Johns Hopkins University

January 26, 2017

## Abstract

This supplementary material contains all the proofs, additional simulation results, and a description of the empirical data.

*Keywords:* bootstrap, confidence band, deconvolution, empirical process theory.

---

\*The authors gratefully acknowledge *please remember to list all relevant funding sources in the unblinded version*

# A Proofs

In what follows, the notation  $\lesssim$  signifies that the left hand side is bounded by the right hand side up to some constant independent of  $n$  and  $x$ .

## A.1 Proof of Theorem 3.1

We first state the following lemmas which will be used in the proof of Theorem 3.1. For a class of measurable functions  $\mathcal{F}$  on a measurable space  $(S, \mathcal{S})$  and a probability measure  $Q$  on  $\mathcal{S}$ , let  $N(\mathcal{F}, \|\cdot\|_{Q,2}, \delta)$  denote the  $\delta$ -covering number for  $\mathcal{F}$  with respect to the  $L^2(Q)$ -seminorm  $\|\cdot\|_{Q,2}$ ; see Section 2.1 in van der Vaart and Wellner (1996) for details.

**Lemma A.1.** *Let  $K$  be a kernel on  $\mathbb{R}$  such that  $\varphi_K$  is supported in  $[-1, 1]$ , and suppose that  $\varphi_\varepsilon$  does not vanish on  $\mathbb{R}$ . Let  $r_n = 1/\inf_{|t| \leq h_n^{-1}} |\varphi_\varepsilon(t)|$ . Consider the class of functions  $\mathcal{K}_n = \{y \mapsto K_n((x - y)/h_n) : x \in \mathbb{R}\}$ , where  $K_n$  denotes the corresponding deconvolution kernel. Then there exist constants  $A, v > 0$  independent of  $n$  such that for all  $n \geq 1$ ,*

$$\sup_Q N(\mathcal{K}_n, \|\cdot\|_{Q,2}, r_n \delta) \leq (A/\delta)^v, \quad 0 < \delta \leq 1,$$

where  $\sup_Q$  is taken over all Borel probability measures  $Q$  on  $\mathbb{R}$ .

In view of Lemma 1 in Giné and Nickl (2009) (see also Proposition 3.6.12 in Giné and Nickl (2016)), Lemma A.1 readily follows if  $K_n$  has quadratic variation  $\lesssim r_n^2$ . Recall that a complex-valued function  $f$  on  $\mathbb{R}$  is said to be of bounded  $p$ -variation for  $p \in [1, \infty)$  if

$$V_p(f) := \sup \left\{ \sum_{\ell=1}^N |f(x_\ell) - f(x_{\ell-1})|^p : -\infty < x_0 < \dots < x_N < \infty, N = 1, 2, \dots \right\}$$

is finite. A function of bounded 2-variation is said to be of bounded quadratic variation. Now, Lemma A.1 follows from the next lemma.

**Lemma A.2.** *Assume the same conditions as in Lemma A.1. Then the deconvolution kernel  $K_n$  is of bounded quadratic variation with  $V_2(K_n) \lesssim r_n^2$ .*

*Proof.* The basic idea of the proof is due to the proof of Lemma 1 in Lounici and Nickl (2011). In view of the continuous embedding of the homogeneous Besov space  $\dot{B}_{2,1}^{1/2}(\mathbb{R})$  into  $BV_2(\mathbb{R})$ , the space of functions of bounded quadratic variation (Bourdaud et al., 2006, Theorem 5), it is enough to show that  $\|K_n\|_{\dot{B}_{2,1}^{1/2}} \lesssim r_n$ , where

$$\|f\|_{\dot{B}_{2,1}^{1/2}} = \int_{\mathbb{R}} \frac{1}{|u|^{3/2}} \left( \int_{\mathbb{R}} |f(x+u) - f(x)|^2 dx \right)^{1/2} du.$$

Let  $\psi_n(t) = \varphi_K(t)/\varphi_\varepsilon(t/h_n)$ , and observe that, using Plancherel's theorem,

$$\int_{\mathbb{R}} |K_n(x+u) - K_n(x)|^2 dx = \frac{1}{2\pi} \int_{\mathbb{R}} |e^{-itu} - 1|^2 |\psi_n(t)|^2 dt = \frac{1}{\pi} \int_{\mathbb{R}} (1 - \cos(tu)) |\psi_n(t)|^2 dt.$$

Using the inequality  $1 - \cos(tu) \leq \min\{2, (tu)^2/2\}$ , we conclude that

$$\begin{aligned} \|K_n\|_{\dot{B}_{2,1}^{1/2}} &\lesssim \left( \int_{\mathbb{R}} t^2 |\psi_n(t)|^2 dt \right)^{1/2} \int_{[-1,1]} \frac{1}{|u|^{1/2}} du + \left( \int_{\mathbb{R}} |\psi_n(t)|^2 dt \right)^{1/2} \int_{[-1,1]^c} \frac{1}{|u|^{3/2}} du \\ &\lesssim r_n. \end{aligned}$$

This completes the proof.  $\square$

**Remark A.1.** Lemma A.1 generalizes a part of Lemma 5.3.5 in Giné and Nickl (2016) that focuses on the case where the bandwidth  $h_n$  takes values in  $\{2^{-k} : k = 1, 2, \dots\}$ . The proof of Lemma A.1 appears to be simpler than that of Lemma 5.3.5 in Giné and Nickl (2016) (but note that Lemma 5.3.5 in Giné and Nickl (2016) also covers wavelet kernels).

**Lemma A.3.** *Assumptions 1–4 imply that, for sufficiently large  $n$ ,  $\inf_{x \in I} \sigma_n^2(x) \gtrsim h_n^{-2\alpha+1}$ .*

*Proof.* The proof is inspired by Fan (1991). Observe first that  $\|K_n\|_{\mathbb{R}} \lesssim h_n^{-\alpha}$ . Second, integration by parts yields that

$$K_n(x) = \frac{1}{2\pi ix} \int_{\mathbb{R}} e^{-itx} \left\{ \frac{\varphi_K(t)}{\varphi_\varepsilon(t/h_n)} \right\}' dt = \frac{1}{2\pi ix} \int_{\mathbb{R}} e^{-itx} \left\{ \frac{\varphi_K'(t)}{\varphi_\varepsilon(t/h_n)} - \frac{\varphi_K(t)\varphi_\varepsilon'(t/h_n)}{h_n\varphi_\varepsilon^2(t/h_n)} \right\} dt.$$

It is not difficult to verify that

$$\int_{\mathbb{R}} \left| \frac{\varphi'_K(t)}{\varphi_\varepsilon(t/h_n)} \right| dt \lesssim h_n^{-\alpha}.$$

Splitting the integral into  $|t/h_n| \leq 1$  and  $|t/h_n| > 1$ , we also see that

$$\left\{ \int_{|t/h_n| \leq 1} + \int_{|t/h_n| > 1} \right\} \left| \frac{\varphi_K(t)\varphi'_\varepsilon(t/h_n)}{h_n\varphi_\varepsilon^2(t/h_n)} \right| dt \lesssim 1 + h_n^{-\alpha} \int_{\mathbb{R}} |t|^{\alpha-1} |\varphi_K(t)| dt,$$

which is  $\lesssim h_n^{-\alpha}$ . This yields that  $h_n^{2\alpha} K_n^2(x) \lesssim 1/x^2$ , and so  $h_n^{2\alpha} K_n^2(x) \lesssim \min\{1, 1/x^2\}$ .

Now, observe that

$$\begin{aligned} |\mathbb{E}[K_n((x-Y)/h_n)]| &= h_n \left| \int_{\mathbb{R}} K(y) f_X(x - h_n y) dy \right| \leq h_n \|f_X\|_{\mathbb{R}} \int_{\mathbb{R}} |K(y)| dy = O(h_n), \text{ and} \\ \mathbb{E}[K_n^2((x-Y)/h_n)] &= h_n \int_{\mathbb{R}} K_n^2(y) f_Y(x - h_n y) dy. \end{aligned}$$

So, it is enough to prove that

$$\inf_{x \in I} \int_{\mathbb{R}} K_n^2(y) f_Y(x - h_n y) dy \gtrsim (1 - o(1)) h_n^{-2\alpha}.$$

To this end, since

$$\int_{\mathbb{R}} K_n^2(y) dy = \frac{1}{2\pi} \int_{\mathbb{R}} \frac{|\varphi_K(t)|^2}{|\varphi_\varepsilon(t/h_n)|^2} dt \gtrsim h_n^{-2\alpha},$$

by Plancherel's theorem, it is enough to prove that as  $n \rightarrow \infty$ ,

$$\sup_{x \in I} \left| h_n^{2\alpha} \int_{\mathbb{R}} K_n^2(y) \{f_Y(x - h_n y) - f_Y(x)\} dy \right| \rightarrow 0.$$

Since  $f_Y$  is continuous and  $I$  is compact, for any  $\rho > 0$ , there exists  $\delta > 0$  such that  $\sup_{x \in I} |f_Y(x - y) - f_Y(x)| \leq \rho$  whenever  $|y| \leq \delta$ . So,

$$\begin{aligned} &\sup_{x \in I} h_n^{2\alpha} \int_{\mathbb{R}} K_n^2(y) |f_Y(x - h_n y) - f_Y(x)| dy \\ &\lesssim \rho \int_{|y| \leq \delta/h_n} \min\{1, 1/y^2\} dy + 2 \|f_Y\|_{\mathbb{R}} \int_{|y| > \delta/h_n} y^{-2} dy \lesssim \rho + o(1), \end{aligned}$$

which yields the desired conclusion.  $\square$

*Proof of Theorem 3.1.* We divide the proof into three steps.

**Step 1.** (Gaussian approximation to  $Z_n^*$ ). Recall the empirical process  $Z_n^*(x), x \in I$  defined as

$$\begin{aligned} Z_n^*(x) &= \frac{\sqrt{n}h_n\{\widehat{f}_X^*(x) - \mathbb{E}[\widehat{f}_X^*(x)]\}}{\sigma_n(x)} \\ &= \frac{1}{\sigma_n(x)\sqrt{n}} \sum_{j=1}^n \{K_n((x - Y_j)/h_n) - \mathbb{E}[K_n((x - Y)/h_n)]\}, \quad x \in I. \end{aligned}$$

Consider the class of functions

$$\mathcal{G}_n = \left\{ \frac{1}{\sigma_n(x)} \{K_n((x - \cdot)/h_n) - \mathbb{E}[K_n((x - Y)/h_n)]\} : x \in I \right\},$$

together with the empirical process indexed by  $\mathcal{G}_n$  defined as

$$\mathcal{G}_n \ni g \mapsto \nu_n(g) = \frac{1}{\sqrt{n}} \sum_{j=1}^n g(Y_j).$$

Observe that  $\|Z_n^*\|_I = \|\nu_n\|_{\mathcal{G}_n}$ . We apply Corollary 2.2 in Chernozhukov et al. (2014a) to  $\|\nu_n\|_{\mathcal{G}_n}$ . First, since the set  $\{1/\sigma_n(x) : x \in I\}$  is bounded with  $\|1/\sigma_n(\cdot)\|_I \lesssim h_n^{\alpha-1/2}$ , in view of Lemma A.1 of the present paper and Corollary A.1 in Chernozhukov et al. (2014a), there exist constants  $A', v' > 0$  independent of  $n$  such that

$$\sup_Q N(\mathcal{G}_n, \|\cdot\|_{Q,2}, \delta/\sqrt{h_n}) \leq (A'/\delta)^{v'}, \quad 0 < \delta \leq 1, \quad (\text{A.1})$$

which ensures that there exists a tight Gaussian random variable  $G_n$  in  $\ell^\infty(\mathcal{G}_n)$  with mean zero and the same covariance function as  $\nu_n$  (cf. Chernozhukov et al., 2014a, Lemma 2.1). Since  $\|K_n((x - \cdot)/h_n)/\sigma_n(x)\|_{\mathbb{R}} \lesssim 1/\sqrt{h_n}$  and  $\text{Var}(K_n((x - Y)/h_n)/\sigma_n(x)) = 1$ , application of Corollary 2.2 in Chernozhukov et al. (2014a) with  $q = \infty, b \lesssim 1/\sqrt{h_n}, \sigma = 1, A \lesssim 1, v \lesssim 1$ , and  $\gamma = 1/\log n$ , yields that there exists a sequence of random variables  $W_n$  with  $W_n \stackrel{d}{=} \|G_n\|_{\mathcal{G}_n}$  (where the notation  $\stackrel{d}{=}$  signifies equality in distribution) and such that

$$\|\nu_n\|_{\mathcal{G}_n} - W_n = O_P\{(\log n)/(nh_n)^{1/6}\}, \quad (\text{A.2})$$

where the left hand side is equal to  $\|Z_n^*\|_I - W_n$ .

Next, for  $f_{n,x}(\cdot) = \{K_n((x - \cdot)/h_n) - \mathbb{E}[K_n((x - Y)/h_n)]\}/\sigma_n(x)$ , let

$$Z_n^G(x) = G_n(f_{n,x}), \quad x \in I,$$

and observe that  $Z_n^G$  is a tight Gaussian random variable in  $\ell^\infty(I)$  with mean zero and the same covariance function as  $Z_n^*$ , and such that  $\|Z_n^G\|_I = \|G_n\|_{\mathcal{G}_n} \stackrel{d}{=} W_n$ . It is worth noting that deducing from (A.2) a bound on

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|Z_n^*\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}|$$

is a non-trivial step, because the distribution of the approximating Gaussian process  $Z_n^G$  changes with  $n$ . To this end, we rely on the anti-concentration inequality for the supremum of a Gaussian process, which yields that

$$\sup_{z \in \mathbb{R}; \Delta > 0} \frac{\mathbb{P}\{z - \Delta \leq \|Z_n^G\|_I \leq z + \Delta\}}{\Delta} \lesssim \mathbb{E}[\|Z_n^G\|_I]. \quad (\text{A.3})$$

See Corollary 2.1 in Chernozhukov et al. (2014b) (see also Theorem 3 in Chernozhukov et al. (2015)). To apply this inequality, we shall bound  $\mathbb{E}[\|Z_n^G\|_I] = \mathbb{E}[\|G_n\|_{\mathcal{G}_n}]$ , but given (A.1) and  $\text{Var}(K_n((x - Y)/h_n)/\sigma_n(x)) = 1$ , Dudley's entropy integral bound (cf. van der Vaart and Wellner, 1996, Corollary 2.2.8) yields that

$$\mathbb{E}[\|G_n\|_{\mathcal{G}_n}] \lesssim \int_0^1 \sqrt{1 + \log(1/(\delta\sqrt{h_n}))} d\delta \lesssim \sqrt{\log(1/h_n)}.$$

Now, combining (A.2) with the anti-concentration inequality (A.3), we conclude that

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|Z_n^*\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \rightarrow 0,$$

provided that  $\frac{(\log n)\sqrt{\log(1/h_n)}}{(nh_n)^{1/6}} \rightarrow 0$ , which is satisfied under our assumption.

**Step 2.** (Gaussian approximation to the intermediate process). Define the intermediate process

$$\tilde{Z}_n(x) = \frac{\sqrt{nh_n}\{\widehat{f}_X(x) - \mathbb{E}[\widehat{f}_X^*(x)]\}}{\sigma_n(x)}, \quad x \in I,$$

where the difference from  $\widehat{Z}_n(x)$  is that  $\widehat{\sigma}_n(x)$  is replaced by  $\sigma_n(x)$ . In this step, we wish to prove that

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\tilde{Z}_n\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \rightarrow 0, \quad (\text{A.4})$$

where  $Z_n^G$  is given in the previous step.

Since  $\varphi_\varepsilon$  does not vanish on  $\mathbb{R}$ , we have that  $\varphi_Y(t) = 0$  if and only if  $\varphi_X(t) = 0$ , from which we have

$$\begin{aligned} \widehat{f}_X^*(x) &= \frac{1}{2\pi} \left\{ \int_{\{\varphi_X \neq 0\}} e^{-itx} \varphi_K(th_n) \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} \varphi_X(t) dt + \int_{\{\varphi_X = 0\}} e^{-itx} \frac{\varphi_K(th_n)}{\varphi_\varepsilon(t)} \widehat{\varphi}_Y(t) dt \right\}, \text{ and} \\ \widehat{f}_X(x) &= \frac{1}{2\pi} \left\{ \int_{\{\varphi_X \neq 0\}} e^{-itx} \varphi_K(th_n) \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} \varphi_X(t) dt + \int_{\{\varphi_X = 0\}} e^{-itx} \frac{\varphi_K(th_n)}{\varphi_\varepsilon(t)} \widehat{\varphi}_Y(t) \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} dt \right\}. \end{aligned}$$

So, letting

$$\tilde{f}_X(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \varphi_K(th_n) \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} \varphi_X(t) dt,$$

we obtain the following decomposition:

$$\begin{aligned} &\widehat{f}_X(x) - \widehat{f}_X^*(x) \\ &= \left[ \{\widehat{f}_X(x) - \tilde{f}_X(x)\} - \{\widehat{f}_X^*(x) - \mathbb{E}[\widehat{f}_X^*(x)]\} \right] + \{\tilde{f}_X(x) - \mathbb{E}[\widehat{f}_X^*(x)]\} \\ &= \frac{1}{2\pi} \int_{\{\varphi_X \neq 0\}} e^{-itx} \varphi_K(th_n) \left\{ \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} - 1 \right\} \left\{ \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right\} \varphi_X(t) dt \\ &\quad + \frac{1}{2\pi} \int_{\{\varphi_X = 0\}} e^{-itx} \frac{\varphi_K(th_n)}{\varphi_\varepsilon(t)} \widehat{\varphi}_Y(t) \left\{ \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right\} dt \\ &\quad + \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \varphi_K(th_n) \left\{ \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right\} \varphi_X(t) dt. \end{aligned}$$

Hence the Cauchy-Schwarz inequality yields that

$$\begin{aligned}
|\widehat{f}_X(x) - \widehat{f}_X^*(x)|^2 &\lesssim \left\{ \int_{\{\varphi_X \neq 0\} \cap [-h_n^{-1}, h_n^{-1}]} \left| \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} - 1 \right|^2 |\varphi_X(t)|^2 dt \right\} \left\{ \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 dt \right\} \\
&\quad + h_n^{-2\alpha} \left\{ \int_{\{\varphi_X = 0\} \cap [-h_n^{-1}, h_n^{-1}]} |\widehat{\varphi}_Y(t)|^2 dt \right\} \left\{ \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 dt \right\} \\
&\quad + \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 |\varphi_X(t)| dt.
\end{aligned}$$

We will bound the following four terms:

$$\begin{aligned}
&\int_{\{\varphi_X \neq 0\} \cap [-h_n^{-1}, h_n^{-1}]} \left| \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} - 1 \right|^2 |\varphi_X(t)|^2 dt, \quad \int_{\{\varphi_X = 0\} \cap [-h_n^{-1}, h_n^{-1}]} |\widehat{\varphi}_Y(t)|^2 dt, \\
&\int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 dt, \quad \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 |\varphi_X(t)| dt.
\end{aligned}$$

To bound the first term, pick and fix any  $t \in \mathbb{R}$  such that  $\varphi_Y(t) \neq 0$ ; let  $\zeta_j = e^{itY_j}/\varphi_Y(t)$  for  $j = 1, \dots, n$ . Then we have

$$\mathbb{E} \left[ \left| \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} - 1 \right|^2 \right] = \mathbb{E} \left[ \left| \frac{1}{n} \sum_{j=1}^n \{\zeta_j - \mathbb{E}[\zeta_j]\} \right|^2 \right] \leq \frac{1}{n|\varphi_Y(t)|^2}.$$

Therefore,

$$\mathbb{E} \left[ \int_{\{\varphi_X \neq 0\} \cap [-h_n^{-1}, h_n^{-1}]} \left| \frac{\widehat{\varphi}_Y(t)}{\varphi_Y(t)} - 1 \right|^2 |\varphi_X(t)|^2 dt \right] \leq n^{-1} \int_{-h_n^{-1}}^{h_n^{-1}} \frac{1}{|\varphi_\varepsilon(t)|^2} dt \lesssim h_n^{-2\alpha} (nh_n)^{-1}.$$

On the other hand, since  $\varphi_Y(t) = 0$  whenever  $\varphi_X(t) = 0$ , we have

$$\mathbb{E} \left[ \int_{\{\varphi_X = 0\} \cap [-h_n^{-1}, h_n^{-1}]} |\widehat{\varphi}_Y(t)|^2 dt \right] \leq 2(nh_n)^{-1}.$$

To bound the third and fourth terms, we first note that, from Lemma B.1 in the supplementary material together with the fact that  $\mathbb{E}[|\varepsilon|^p] < \infty$  for some  $p > 0$ ,

$$\|\widehat{\varphi}_\varepsilon - \varphi_\varepsilon\|_{[-h_n^{-1}, h_n^{-1}]} = O_P\{m^{-1/2} \log(1/h_n)\}, \tag{A.5}$$

which is  $o_{\mathbb{P}}(h_n^\alpha)$  by Assumption 6 (b). Hence

$$\inf_{t \in [-h_n^{-1}, h_n^{-1}]} |\widehat{\varphi}_\varepsilon(t)| \geq \inf_{t \in [-h_n^{-1}, h_n^{-1}]} |\varphi_\varepsilon(t)| - o_{\mathbb{P}}(h_n^\alpha) \gtrsim (1 - o_{\mathbb{P}}(1))h_n^\alpha,$$

from which we have

$$\begin{aligned} \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 dt &\lesssim h_n^{-2\alpha} \int_{-h_n^{-1}}^{h_n^{-1}} |\varphi_\varepsilon(t) - \widehat{\varphi}_\varepsilon(t)|^2 dt = O_{\mathbb{P}}(h_n^{-2\alpha-1}m^{-1}), \\ \int_{-h_n^{-1}}^{h_n^{-1}} \left| \frac{\varphi_\varepsilon(t)}{\widehat{\varphi}_\varepsilon(t)} - 1 \right|^2 |\varphi_X(t)| dt &= O_{\mathbb{P}}(h_n^{-2\alpha}m^{-1}), \end{aligned}$$

where we have used the fact that  $|\varphi_X|$  is integrable on  $\mathbb{R}$ .

Taking these together, we have

$$\|\widehat{f}_X - \widehat{f}_X^*\|_{\mathbb{R}} = o_{\mathbb{P}} \left[ h_n^{-\alpha} \{(nh_n) \log(1/h_n)\}^{-1/2} \right]$$

by Assumption 6 (b), from which we conclude that

$$\|\widetilde{Z}_n - Z_n^*\|_I = o_{\mathbb{P}}\{(\log(1/h_n))^{-1/2}\}.$$

This shows that there exists a sequence  $\Delta_n \downarrow 0$  such that

$$\mathbb{P}\{\|\widetilde{Z}_n - Z_n^*\|_I > \Delta_n (\log(1/h_n))^{-1/2}\} \leq \Delta_n$$

(which follows from the fact that convergence in probability is metrized by the Ky Fan metric; see Theorem 9.2.2 in Dudley (2002)), and so

$$\begin{aligned} \mathbb{P}\{\|\widetilde{Z}_n\|_I \leq z\} &\leq \mathbb{P}\{\|Z_n^*\|_I \leq z + \Delta_n (\log(1/h_n))^{-1/2}\} + \Delta_n \\ &\leq \mathbb{P}\{\|Z_n^G\|_I \leq z + \Delta_n (\log(1/h_n))^{-1/2}\} + o(1) \leq \mathbb{P}\{\|Z_n^G\|_I \leq z\} + o(1) \end{aligned}$$

uniformly in  $z \in \mathbb{R}$ , where the second inequality follows from the previous step, and the last inequality follows from the anti-concentration inequality (A.3). Likewise, we have

$P\{\|\tilde{Z}_n\|_I \leq z\} \geq P\{\|Z_n^G\|_I \leq z\} - o(1)$  uniformly in  $z \in \mathbb{R}$ , so that we obtain the conclusion of this step.

**Step 3.** (Proof of the theorem). Observe that

$$\begin{aligned} \|\hat{K}_n - K_n\|_{\mathbb{R}} &\lesssim \int_{\mathbb{R}} \left| \frac{1}{\hat{\varphi}_\varepsilon(t/h_n)} - \frac{1}{\varphi_\varepsilon(t/h_n)} \right| |\varphi_K(t)| dt \\ &\leq O_P(h_n^{-2\alpha}) \int_{\mathbb{R}} |\hat{\varphi}_\varepsilon(t/h_n) - \varphi_\varepsilon(t/h_n)| |\varphi_K(t)| dt, \end{aligned}$$

which is  $O_P(h_n^{-2\alpha}m^{-1/2})$  since  $\int_{\mathbb{R}} E[|\hat{\varphi}_\varepsilon(t/h_n) - \varphi_\varepsilon(t/h_n)|] |\varphi_K(t)| dt \lesssim m^{-1/2}$ . Together with the fact that  $\|K_n\|_{\mathbb{R}} = O(h_n^{-\alpha})$ , we have

$$\|\hat{K}_n^2 - K_n^2\|_{\mathbb{R}} \leq \|\hat{K}_n - K_n\|_{\mathbb{R}} \|\hat{K}_n + K_n\|_{\mathbb{R}} = O(h_n^{-3\alpha}m^{-1/2}),$$

which yields that

$$\hat{\sigma}_n^2(x) = \frac{1}{n} \sum_{j=1}^n K_n^2((x - Y_j)/h_n) - \left( \frac{1}{n} \sum_{j=1}^n K_n((x - Y_j)/h_n) \right)^2 + O_P(h_n^{-3\alpha}m^{-1/2})$$

uniformly in  $x \in I$ . It is not difficult to verify that

$$\begin{aligned} \frac{1}{n} \sum_{j=1}^n K_n^2((x - Y_j)/h_n) &= E[K_n^2((x - Y)/h_n)] + O_P(h_n^{-2\alpha}n^{-1/2}), \text{ and} \\ \frac{1}{n} \sum_{j=1}^n K_n((x - Y_j)/h_n) &= E[K_n((x - Y)/h_n)] + O_P(h_n^{-\alpha}n^{-1/2}) \end{aligned}$$

uniformly in  $x \in I$ . Indeed, in view of Lemma A.1 of the present paper and Corollary A.1 in Chernozhukov et al. (2014a), these estimates follow from Theorem 2.14.1 in van der Vaart and Wellner (1996). Therefore, we have  $\hat{\sigma}_n^2(x) = \sigma_n^2(x) + O_P\{h_n^{-2\alpha}(h_n^{-\alpha}m^{-1/2} + n^{-1/2})\}$  uniformly in  $x \in I$ , and so  $\hat{\sigma}_n^2(x)/\sigma_n^2(x) = 1 + O_P\{h_n^{-1}(h_n^{-\alpha}m^{-1/2} + n^{-1/2})\}$  uniformly in  $x \in I$  by Lemma A.3, where  $h_n^{-1}(h_n^{-\alpha}m^{-1/2} + n^{-1/2}) = o\{(\log(1/h_n))^{-1}\}$  by Assumption 6. This yields that

$$\|\sigma_n(\cdot)/\hat{\sigma}_n(\cdot) - 1\|_I = o_P\{(\log(1/h_n))^{-1}\}.$$

By Steps 1 and 2 together with the fact that  $E[\|Z_n^G\|_I] = O(\sqrt{\log(1/h_n)})$ , we see that  $\|\tilde{Z}_n\|_I = O_P(\sqrt{\log(1/h_n)})$ . So,

$$\|\hat{Z}_n - \tilde{Z}_n\|_I \leq \|\sigma_n(\cdot)/\hat{\sigma}_n(\cdot) - 1\|_I \|\tilde{Z}_n\|_I = o_P\{(\log(1/h_n))^{-1/2}\},$$

and arguing as in the last part of the proof of Step 2, we conclude that

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\hat{Z}_n\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \rightarrow 0.$$

This completes the proof of Theorem 3.1.  $\square$

## A.2 Proofs of Corollaries 3.1 and 3.2

*Proof of Corollary 3.1.* Step 2 in the proof of Theorem 3.1 yields that  $\|\hat{f}_X - \hat{f}_X^*\|_{\mathbb{R}} = o_P\{h_n^{-\alpha}(nh_n)^{-1/2}\}$  (it is not difficult to verify that Assumption 4 was not used to derive this rate). Hence we have to show that  $\|\hat{f}_X^*(\cdot) - E[\hat{f}_X^*(\cdot)]\|_{\mathbb{R}} = O_P\{h_n^{-\alpha}(nh_n)^{-1/2}\sqrt{\log(1/h_n)}\}$ . To this end, we make use of Corollary 5.1 in Chernozhukov et al. (2014a). Invoke Lemma A.1 and observe that  $\|K_n\|_{\mathbb{R}} \lesssim h_n^{-\alpha}$  and  $\sigma_n^2(x) \lesssim h_n^{-2\alpha+1}$ . Application of Corollary 5.1 in Chernozhukov et al. (2014a) to  $\mathcal{K}_n$  then yields that

$$\begin{aligned} & \mathbb{E} \left[ \left\| \sum_{j=1}^n \{K((\cdot - Y_j)/h_n) - E[K_n((\cdot - Y)/h_n)]\} \right\|_{\mathbb{R}} \right] \\ & \lesssim h_n^{-\alpha} \sqrt{nh_n \log(1/h_n)} + h_n^{-\alpha} \log(1/h_n) \\ & \lesssim h_n^{-\alpha} \sqrt{nh_n \log(1/h_n)}, \end{aligned}$$

which in turn yields that

$$\mathbb{E}[\|\hat{f}_X^*(\cdot) - E[\hat{f}_X^*(\cdot)]\|_{\mathbb{R}}] \lesssim h_n^{-\alpha}(nh_n)^{-1/2}\sqrt{\log(1/h_n)}.$$

This completes the proof.  $\square$

*Proof of Corollary 3.2.* Since  $\varphi_K$  is  $(k + 3)$ -times continuously differentiable, the kernel  $K(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-itx} \varphi_K(t) dt$  is such that  $\int_{\mathbb{R}} |x|^{k+1} |K(x)| dx < \infty$  and  $\int_{\mathbb{R}} x^\ell K(x) dx = i^{-\ell} \varphi_K^{(\ell)}(0) = 0$  for all  $\ell = 1, \dots, k$ . Hence

$$\begin{aligned} \|\mathbb{E}[\widehat{f}_X^*(\cdot)] - f_X(\cdot)\|_{\mathbb{R}} &= \left\| \int_{\mathbb{R}} \{f_X(\cdot - h_n y) - f_X(\cdot)\} K(y) dy \right\|_{\mathbb{R}} \\ &\leq \frac{B h_n^\beta}{k!} \int_{\mathbb{R}} |y|^\beta |K(y)| dy. \end{aligned}$$

Combining the result of Corollary 3.1, and in view of the remark after Assumption 6, we obtain the desired conclusion.  $\square$

### A.3 Proofs of Theorem 3.2 and Corollary 3.3

*Proof of Theorem 3.2.* We divide the proof into three steps.

**Step 1.** Define

$$Z_n^\xi(x) = \frac{1}{\sigma_n(x) \sqrt{n}} \sum_{j=1}^n \xi_j \left\{ K_n((x - Y_j)/h_n) - n^{-1} \sum_{j'=1}^n K_n((x - Y_{j'})/h_n) \right\}$$

for  $x \in I$ . We first prove that

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|Z_n^\xi\|_I \leq z \mid \mathcal{D}_n\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \xrightarrow{\mathbb{P}} 0.$$

To this end, we make use of Theorem 2.2 in Chernozhukov et al. (2016). Recall the class of functions  $\mathcal{G}_n$  defined in the proof of Theorem 3.1, and let

$$\nu_n^\xi(g) = \frac{1}{\sqrt{n}} \sum_{j=1}^n \xi_j \left\{ g(Y_j) - n^{-1} \sum_{j'=1}^n g(Y_{j'}) \right\}, \quad g \in \mathcal{G}_n.$$

Then application of Theorem 2.2 in Chernozhukov et al. (2016) to  $\|\nu_n\|_{\mathcal{G}_n}$  with  $B(f) \equiv 0, b \lesssim 1/\sqrt{h_n}, \sigma = 1, A \lesssim 1, v \lesssim 1, \gamma = 1/\log n$ , and  $q$  sufficiently large, yields that there

exists a random variable  $W_n^\xi$  of which the conditional distribution given  $\mathcal{D}_n$  is the same as the distribution of  $\|G_n\|_{\mathcal{G}_n}$  ( $= \|Z_n^G\|_I$ ), and such that

$$\|\nu_n^\xi\|_{\mathcal{G}_n} - W_n^\xi = o_{\mathbb{P}}\{(\log(1/h_n))^{-1/2}\},$$

which shows that there exists a sequence  $\Delta_n \downarrow 0$  such that

$$\mathbb{P}\{\|\nu_n^\xi\|_{\mathcal{G}_n} - W_n^\xi > \Delta_n(\log(1/h_n))^{-1/2} \mid \mathcal{D}_n\} \xrightarrow{\mathbb{P}} 0$$

by Markov's inequality. Since  $\|\nu_n^\xi\|_{\mathcal{G}_n} = \|Z_n^\xi\|_I$ , we have

$$\begin{aligned} \mathbb{P}\{\|Z_n^\xi\|_I \leq z \mid \mathcal{D}_n\} &\leq \mathbb{P}\{W_n^\xi \leq z + \Delta_n(\log(1/h_n))^{-1/2} \mid \mathcal{D}_n\} + o_{\mathbb{P}}(1) \\ &= \mathbb{P}\{\|Z_n^G\|_I \leq z + \Delta_n(\log(1/h_n))^{-1/2}\} + o_{\mathbb{P}}(1) \end{aligned}$$

uniformly in  $z \in \mathbb{R}$ , and the anti-concentration inequality (A.3) yields that

$$\mathbb{P}\{\|Z_n^G\|_I \leq z + \Delta_n(\log(1/h_n))^{-1/2}\} \leq \mathbb{P}\{\|Z_n^G\|_I \leq z\} + o(1)$$

uniformly in  $z \in \mathbb{R}$ . Likewise, we have

$$\mathbb{P}\{\|Z_n^\xi\|_I \leq z \mid \mathcal{D}_n\} \geq \mathbb{P}\{\|Z_n^G\|_I \leq z\} - o_{\mathbb{P}}(1)$$

uniformly in  $z \in \mathbb{R}$ . Therefore, we obtain the conclusion of this step.

**Step 2.** In view of the proof of Step 1, in order to prove the result (3.2), it is enough to prove that

$$\|\widehat{Z}_n^\xi - Z_n^\xi\|_I = o_{\mathbb{P}}\{(\log(1/h_n))^{-1/2}\}.$$

Define

$$\widetilde{Z}_n^\xi = \frac{1}{\sigma_n(x)\sqrt{n}} \sum_{j=1}^n \xi_j \left\{ \widehat{K}_n((x - Y_j)/h_n) - n^{-1} \sum_{j'=1}^n \widehat{K}_n((x - Y_{j'})/h_n) \right\}$$

for  $x \in I$ . We first prove that  $\|\tilde{Z}_n^\xi - Z_n^\xi\|_I = o_P\{(\log(1/h_n))^{-1/2}\}$ . Step 2 in the proof of Theorem 3.1 shows that

$$\left\| \frac{1}{\sigma_n(\cdot)\sqrt{n}} \sum_{j=1}^n \{\widehat{K}_n((\cdot - Y_j)/h_n) - K_n((\cdot - Y_j)/h_n)\} \right\|_I$$

is  $o_P\{(\log(1/h_n))^{-1/2}\}$  (which is more than what we need), and so it remains to prove that

$$\left\| \frac{1}{\sigma_n(\cdot)\sqrt{n}} \sum_{j=1}^n \xi_j \{\widehat{K}_n((\cdot - Y_j)/h_n) - K_n((\cdot - Y_j)/h_n)\} \right\|_I$$

is  $o_P\{(\log(1/h_n))^{-1/2}\}$ . Since  $\|1/\sigma_n(\cdot)\|_I \lesssim h_n^{\alpha-1/2}$ , it is enough to prove that

$$\left\| \sum_{j=1}^n \xi_j \{\widehat{K}_n((\cdot - Y_j)/h_n) - K_n((\cdot - Y_j)/h_n)\} \right\|_I$$

is  $o_P\{h_n^{-\alpha}(nh_n)^{1/2}(\log(1/h_n))^{-1/2}\}$ . Observe that

$$\begin{aligned} & \left| \sum_{j=1}^n \xi_j \{\widehat{K}_n((x - Y_j)/h_n) - K_n((x - Y_j)/h_n)\} \right| \\ & \lesssim \int_{\mathbb{R}} \left| \sum_{j=1}^n \xi_j e^{itY_j/h_n} \right| \left| \frac{1}{\widehat{\varphi}_\varepsilon(t/h_n)} - \frac{1}{\varphi_\varepsilon(t/h_n)} \right| |\varphi_K(t)| dt \\ & \lesssim \left\{ \int_{\mathbb{R}} \left| \sum_{j=1}^n \xi_j e^{itY_j/h_n} \right|^2 |\varphi_K(t)| dt \right\}^{1/2} \left\{ \int_{\mathbb{R}} \left| \frac{1}{\widehat{\varphi}_\varepsilon(t/h_n)} - \frac{1}{\varphi_\varepsilon(t/h_n)} \right|^2 |\varphi_K(t)| dt \right\}^{1/2} \\ & = O_P(n^{1/2}h_n^{-2\alpha}m^{-1/2}), \end{aligned}$$

which is  $o_P\{h_n^{-\alpha}(nh_n)^{1/2}(\log(1/h_n))^{-1/2}\}$  by Assumption 6 (b). Therefore, we have  $\|\tilde{Z}_n^\xi - Z_n^\xi\|_I = o_P\{(\log(1/h_n))^{-1/2}\}$ .

By Step 1, the fact that  $E[\|Z_n^G\|_I] = O(\sqrt{\log(1/h_n)})$  (see the proof of Theorem 3.1), and the previous result, we see that  $\|\tilde{Z}_n^\xi\|_I = O_P(\sqrt{\log(1/h_n)})$ . Now, because  $\|\sigma_n(\cdot)/\widehat{\sigma}_n(\cdot) -$

$1\|_I = o_{\mathbb{P}}\{(\log(1/h_n))^{-1}\}$  from Step 3 in the proof of Theorem 3.1, we conclude that

$$\|\widehat{Z}_n^\xi - \widetilde{Z}_n^\xi\|_I \leq \|\sigma_n(\cdot)/\widehat{\sigma}_n(\cdot) - 1\|_I \|\widetilde{Z}_n^\xi\|_I = o_{\mathbb{P}}\{(\log(1/h_n))^{-1/2}\},$$

which leads to (3.2).

**Step 3.** In this step, we shall verify the last two assertions of the theorem. The result (3.2) implies that there exists a sequence  $\Delta_n \downarrow 0$  such that with probability greater than  $1 - \Delta_n$ ,

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\widehat{Z}_n^\xi\|_I \leq z \mid \mathcal{D}_n\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \leq \Delta_n. \quad (\text{A.6})$$

Further, taking  $\Delta_n \downarrow 0$  more slowly if necessary, we have

$$\sup_{z \in \mathbb{R}} |\mathbb{P}\{\|\widehat{Z}_n\|_I \leq z\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\}| \leq \Delta_n.$$

Let  $\mathcal{E}_n$  denote the event on which (A.6) happens, and let  $c_n^G(u)$  denote the  $u$ -quantile of  $\|Z_n^G\|_I$  for  $u \in (0, 1)$ . Then on the event  $\mathcal{E}_n$ ,

$$\mathbb{P}\{\|\widehat{Z}_n^\xi\|_I \leq c_n^G(1 - \tau + \Delta_n) \mid \mathcal{D}_n\} \geq \mathbb{P}\{\|Z_n^G\|_I \leq c_n^G(1 - \tau + \Delta_n)\} - \Delta_n = 1 - \tau,$$

where the last equality holds since  $\|Z_n^G\|_I$  has a continuous distribution function (recall the anti-concentration inequality (A.3)). This yields that the inequality

$$\widehat{c}_n(1 - \tau) \leq c_n^G(1 - \tau + \Delta_n)$$

holds on  $\mathcal{E}_n$ , and so

$$\begin{aligned} \mathbb{P}\{\|\widehat{Z}_n\|_I \leq \widehat{c}_n(1 - \tau)\} &\leq \mathbb{P}\{\|\widehat{Z}_n\|_I \leq c_n^G(1 - \tau + \Delta_n)\} + \mathbb{P}(\mathcal{E}_n^c) \\ &\leq \mathbb{P}\{\|Z_n^G\|_I \leq c_n^G(1 - \tau + \Delta_n)\} + 2\Delta_n \\ &= 1 - \tau + 3\Delta_n. \end{aligned}$$

Likewise, we have

$$\mathbb{P}\{\|\widehat{Z}_n\|_I \leq \widehat{c}_n(1 - \tau)\} \geq 1 - \tau - 3\Delta_n.$$

This leads to the result (3.3).

Finally, the Borell-Sudakov-Tsirelson inequality (van der Vaart and Wellner, 1996, Lemma A.2.2) yields that

$$c_n^G(1 - \tau + \Delta_n) \leq \mathbb{E}[\|Z_n^G\|_I] + \sqrt{2 \log(1/(\tau - \Delta_n))} \lesssim \sqrt{\log(1/h_n)},$$

which implies that  $\widehat{c}_n(1 - \tau) = O_{\mathbb{P}}(\sqrt{\log(1/h_n)})$ . Furthermore,

$$\sup_{x \in I} \widehat{\sigma}_n(x) \leq \sup_{x \in I} \sigma_n(x) \cdot \sup_{x \in I} \frac{\widehat{\sigma}_n(x)}{\sigma_n(x)} = O_{\mathbb{P}}(h_n^{-\alpha+1/2}).$$

Therefore, we have

$$\sup_{x \in I} s_n(x) = 2 \sup_{x \in I} \frac{\widehat{\sigma}_n(x)}{\sqrt{nh_n}} \widehat{c}_n(1 - \tau) = O_{\mathbb{P}} \left\{ h_n^{-\alpha} (nh_n)^{-1/2} \sqrt{\log(1/h_n)} \right\}.$$

This completes the proof.  $\square$

*Proof of Corollary 3.3.* Recall the stochastic process  $\widetilde{Z}_n(x), x \in I$  defined in the proof of Theorem 3.1. Observe that  $\|f_X(\cdot) - \mathbb{E}[\widehat{f}_X^*(\cdot)]\|_{\mathbb{R}} = O(h_n^\beta)$ . Condition (3.4) then yields that

$$\begin{aligned} \frac{\sqrt{nh_n}(\widehat{f}_X(x) - f_X(x))}{\widehat{\sigma}_n(x)} &= \{1 + o_{\mathbb{P}}\{(\log(1/h_n))^{-1}\}\} \frac{\sqrt{nh_n}(\widehat{f}_X(x) - f_X(x))}{\sigma_n(x)} \\ &= \{1 + o_{\mathbb{P}}\{(\log(1/h_n))^{-1}\}\} \widetilde{Z}_n(x) + o\{(\log(1/h_n))^{-1/2}\} \\ &= \widetilde{Z}_n(x) + o_{\mathbb{P}}\{(\log(1/h_n))^{-1/2}\} \end{aligned}$$

uniformly in  $x \in I$ , where we have used the facts that  $\|\sigma_n(\cdot)/\widehat{\sigma}_n(\cdot) - 1\|_I = o_{\mathbb{P}}\{(\log(1/h_n))^{-1}\}$  and  $\|\widetilde{Z}_n\|_I = O_{\mathbb{P}}(\sqrt{\log(1/h_n)})$  (these estimates are derived in the proof of Theorem 3.1).

Using the anti-concentration inequality (A.3) together with the result of Step 2 in the proof of Theorem 3.1, we have

$$\sup_{z \in \mathbb{R}} \left| \mathbb{P} \left\{ \left\| \frac{\sqrt{nh_n}(\widehat{f}_X(\cdot) - f_X(\cdot))}{\widehat{\sigma}_n(\cdot)} \right\|_I \leq z \right\} - \mathbb{P}\{\|Z_n^G\|_I \leq z\} \right| \rightarrow 0.$$

Now, arguing as in Step 3 in the proof of Theorem 3.2, we conclude that

$$\mathbb{P} \left\{ \left\| \frac{\sqrt{nh_n}(\widehat{f}_X(\cdot) - f_X(\cdot))}{\widehat{\sigma}_n(\cdot)} \right\|_I \leq \widehat{c}_n(1 - \tau) \right\} \rightarrow 1 - \tau,$$

which yields the desired result.  $\square$

## A.4 Proofs for Section 6

The proof of Theorem 6.1 is almost identical to the proofs of Theorems 3.1 and 3.2 in the ordinary smooth case. The only changes that have to be taken into account are (6.1) and (6.2), which imply that  $\|K_n((x - \cdot)/h_n)/\sigma_n(x)\|_{\mathbb{R}} \lesssim h_n^{-\gamma(1+\lambda)}$ , for example. To avoid repetitions, we omit the details for brevity. In view of the proof of Corollary 3.3, Corollary 6.1 directly follows from Theorem 6.1.  $\square$

## B Uniform convergence rates of the empirical characteristic function

In this appendix, we establish rates of convergence of the empirical characteristic function on expanding sets. The proof of the following lemma is due essentially to Neumann and Reiß (2009, Theorem 4.1).

Let  $F$  be a distribution function on  $\mathbb{R}$  with characteristic function  $\varphi(t) = \int_{\mathbb{R}} e^{itx} dF(x)$ , and let  $X_1, \dots, X_n$  be an independent sample from  $F$ . Let  $F_n(x) = n^{-1} \sum_{j=1}^n 1_{(-\infty, x]}(X_j)$

be the empirical distribution function, and let  $\varphi_n(t) = \int_{\mathbb{R}} e^{itx} dF_n(x) = n^{-1} \sum_{j=1}^n e^{itX_j}$  be the empirical characteristic function.

**Lemma B.1.** *Suppose that  $\int_{\mathbb{R}} |x|^p dF(x) < \infty$  for some  $p > 0$ . Then for any  $\delta > 0$  and any  $T_n \rightarrow \infty$ , we have*

$$\|\varphi_n - \varphi\|_{[-T_n, T_n]} = O_{\mathbb{P}}\{n^{-1/2}(\log T_n)^{1/2+\delta}\}.$$

*Proof.* Let  $w(t) = (\log(e + |t|))^{-1/2-\delta}$ . According to Theorem 4.1 in Neumann and Reiß (2009), it follows that

$$C := \sup_{n \geq 1} \mathbb{E}[\|\sqrt{n}(\varphi_n - \varphi)w\|_{\mathbb{R}}] < \infty.$$

Now, because

$$\|\sqrt{n}(\varphi_n - \varphi)w\|_{\mathbb{R}} \geq \sqrt{n}\|\varphi_n - \varphi\|_{[-T_n, T_n]} \inf_{|t| \leq T_n} w(t),$$

we conclude that

$$\mathbb{E}[\|\varphi_n - \varphi\|_{[-T_n, T_n]}] \leq \frac{C}{\sqrt{n} \inf_{|t| \leq T_n} w(t)} = O\{n^{-1/2}(\log T_n)^{1/2+\delta}\},$$

which leads to the desired result by Markov's inequality.  $\square$

It is worthwhile to point out that the restriction to the set  $|t| \leq T_n$  in Lemma B.1 is essential. In fact, although the class of functions  $\{x \mapsto e^{itx} : t \in \mathbb{R}\}$  is uniformly bounded, it is in general not Glivenko-Cantelli (nor Donsker, of course). See Feurerverger and Mureika (1977) for details.

## C Additional simulation results

In addition to the simulation studies on coverage probabilities presented in the main text, we also use Monte Carlo simulations to show the power of uniform specification tests based on

our uniform confidence band. In the main text, simulation results for coverage probabilities are reported for the true density function  $f_X$ . We now consider a list of alternative specifications of  $f_X$  given by  $f_{X,\mu_X}(x) = (2\pi)^{-1/2}e^{-(x-\mu_X)^2/2}$  for  $\mu_X \in \{0.0, 0.1, 0.2, 0.3, 0.4, 0.5\}$ , and likewise consider a list of alternative specifications given by  $f_{X,\sigma_X}(x) = (2\pi\sigma_X^2)^{-1/2}e^{-x^2/(2\sigma_X^2)}$  for  $\sigma_X \in \{1.0, 1.1, 1.2, 1.3, 1.4, 1.5\}$ . For the errors, we again consider the independent Laplace random vector  $(\varepsilon^{(1)}, \varepsilon^{(2)})$  as in Model 1. Figure 1 plots simulated coverage probabilities for the list of the alternative specifications of  $f_{X,\mu_X}$  (top) and for the list of the alternative specifications of  $f_{X,\sigma_X}$  (bottom) for the nominal coverage probability of  $(1 - \tau) = 0.90$ . The three curves are drawn for each of the three sample sizes  $n \in \{250, 500, 1,000\}$ . Observe that, under the true specification (i.e.,  $\mu_X = 0.0$  in the top graph and  $\sigma_X = 1.0$  in the bottom graph), the simulated coverage probabilities are close to the nominal coverage probability of 0.90, with the case of  $n = 1,000$  being the closest and the case of  $n = 250$  being the farthest. On the other hand, as the specification deviates away from the truth (i.e., as  $\mu_X$  or  $\sigma_X$  increases), the nominal coverage probabilities decrease, with the case of  $n = 1,000$  being the fastest and the case of  $n = 250$  being the slowest. These results evidence the power of the uniform specification tests.

## D Auction Data

The source data for our empirical application can be obtained from the Center for the Study of Auctions, Procurements and Competition Policy hosted by Penn State University. We pre-process bid values in this source data and obtain firms' values  $(Y^{(1)}, Y^{(2)})$  based on an equilibrium restriction (Bayesian Nash equilibrium) for the first-price sealed-bid auction mechanism – we use the same procedure as the one used in Li et al. (2000). See also Guerre et al. (2000). While the original sample consists of 217 tracts with two firms in




Figure 1: Simulated uniform coverage probabilities for alternative specifications of  $f_X$  by estimated confidence bands in  $I = [-2, 2]$  under the standard normal random variable  $X$  and an independent Laplace random vector  $(\varepsilon^{(1)}, \varepsilon^{(2)})$ . For the top graph, the list of alternative specifications are given by  $f_{X, \mu_X}(x) = (2\pi)^{-1/2} e^{-(x-\mu_X)^2/2}$  for  $\mu_X \in \{0.0, 0.1, 0.2, 0.3, 0.4, 0.5\}$ . For the bottom graph, the list of alternative specifications are given by  $f_{X, \sigma_X}(x) = (2\pi\sigma_X^2)^{-1/2} e^{-x^2/(2\sigma_X^2)}$  for  $\sigma_X \in \{1.0, 1.1, 1.2, 1.3, 1.4, 1.5\}$ . The simulated probabilities are computed for the nominal coverage probability of 90% based on 2,000 Monte Carlo iterations.

each tract, we obtain 169 tracts with 2 firms in each tract as a result of trimming. Figure 2 depicts a simple kernel density estimate of the values  $(Y^{(1)}, Y^{(2)})$  in the logarithm of US dollars per acre. This figure essentially reproduces Figure 3 of Li et al. (2000). Note that the value distribution is bimodal.




Figure 2: A kernel density estimate of the values  $(Y^{(1)}, Y^{(2)})$  in the logarithm of US dollars per acre.

## References

- Bourdaud, G., Lanza de Cristoforis, M., and Sickel, W. (2006). Superposition operators and functions of bounded  $p$ -variation. *Rev. Mat. Iberoamericana* **22** 455-487.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014a). Gaussian approximation of suprema of empirical processes. *Ann. Statist.* **42** 1564-1597.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2014b). Anti-concentration and honest, adaptive confidence bands. *Ann. Statist.* **42** 1787-1818.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2015). Comparison and anti-concentration bounds for maxima of Gaussian random vectors. *Probab. Theory Related Fields* **162** 47-70.
- Chernozhukov, V., Chetverikov, D., and Kato, K. (2016). Empirical and multiplier bootstraps for suprema of empirical processes of increasing complexity, and related Gaussian couplings. *Stochastic Process. Appl.*, to appear. arXiv:1502:00352.
- Dudley, R.M. (2002). *Real Analysis and Probability*. Cambridge University Press.
- Fan, J. (1991). Asymptotic normality for deconvolution kernel density estimators. *Sankhya A* **53** 97-110.
- Giné, E. and Nickl, R. (2009). Uniform limit theorems for wavelet density estimators. *Ann. Probab.* **37** 1605-1646.
- Giné, E. and Nickl, R. (2016). *Mathematical Foundations of Infinite-Dimensional Statistical Models*. Cambridge University Press.
- Guerre, E., Perrigne, I., and Vuong, Q. (2000). Optimal nonparametric estimation of first-price auctions. *Econometrica*, **68** 525-574.

- Feurerverger, A. and Mureika, R.A. (1977). Empirical characteristic function and its applications. *Ann. Statist.* **5** 88-97.
- Li, T., Perrigne, I., and Vuong, Q. (2000). Conditionally independent private information in OCS wildcat auctions. *J. Econometrics* **98** 129-161.
- Lounici, K. and Nickl, R. (2011). Global uniform risk bounds for wavelet deconvolution estimators. *Ann. Statist.* **39** 201-231.
- Neumann, M.H. and Reiß, M. (2009). Nonparametric estimation for Lévy processes from low-frequency observations. *Bernoulli* **15** 223-248.
- van der Vaart, A.W. and Wellner, J.A. (1996). *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer.