

# A Nonlinear Panel Unit Root Test under Cross Section Dependence \*

**Mario Cerrato** †

Department of Economics  
University of Glasgow, UK

**Christian de Peretti**

SAF (EA2429), Institute of Financial and Insurance Sciences,  
University of Lyon 1, France

DEFI (EA4265), Department of Economics,  
University of Aix-Marseille 2, France

**Rolf Larsson**

Department of Information Science  
Uppsala University, Sweden

**Nicholas Sarantis**

Centre for International Capital Markets  
London Metropolitan Business School  
London Metropolitan University, UK

July 8, 2009

## **Abstract**

We propose a nonlinear heterogeneous panel unit root test for testing the null hypothesis of unit-roots processes against the alternative that allows a proportion of units to be generated by globally stationary ESTAR processes and a remaining non-zero proportion to be generated by unit root processes. The proposed test is simple to implement and accommodates cross sectional dependence. We show that

---

\*We wish to acknowledge Karim Abadir and Ron Smith for their useful comments and Alexander Kadow for brilliant research assistance on the final version. We also wish to express our gratitude for comments received from the conference participants of 2008 ESEM, 27th–31st August 2008, Milan, Italy; “Inference and Tests in Econometrics: A Tribute to Russel Davidson”, GREQAM, Marseilles, France, 25–26 April 2008; 2008 Scottish Economic Society Annual Conference, Perth, 21st–23rd April 2008; and 2008 Studies in Nonlinear Dynamics and Econometrics (SNDE2008) conference, Federal Reserve Bank, San Francisco, 3rd–4th April 2008.

†Corresponding author: Dr Mario Cerrato, Department of Economics, Adam Smith Building, University of Glasgow, Glasgow G12 8RT, Scotland, UNITED KINGDOM. Email: [m.cerrato@lbs.gla.ac.uk](mailto:m.cerrato@lbs.gla.ac.uk), Tel: +44 (0)141 330 4618, Fax: +44 (0)141 330 4940, web: <http://www.gla.ac.uk/departments/economics>.

the distribution of the test statistic is free of nuisance parameters as  $(N, T) \rightarrow \infty$ . Monte Carlo simulation shows that our test holds correct size and under the hypothesis that data are generated by globally stationary ESTAR processes has a better power than the recent test proposed in Pesaran [2007]. An application to a panel of bilateral real exchange rate series with the US Dollar from the 20 major OECD countries is provided.

**J.E.L. Classification:** C12, C15, C22, C23; F31.

**Keywords:** Nonlinear panel unit root tests, cross sectional dependence.

## 1 Introduction

There is now a large literature on testing for the presence of unit roots in economic and financial variables employing a variety of time series and panel tests<sup>1</sup>. The growth in that area is mainly due to empirical applications on, for example, *Purchasing Power Parity* (PPP) and Growth (see Cerrato and Sarantis [2007a], Emerson and Kao [2006] amongst others).

A weakness of the existing univariate and panel unit root tests is that they are based on the assumption that the underlying variable follows a linear process. However economic theory suggests that generally financial economic variables exhibit nonlinear behaviour. For example, a number of theoretical models in international macroeconomics formalise the notion of nonlinear exchange rate behaviour due to transaction costs (*e.g.* Dumas [1992], Sercu and Uppal [1995], O’Connell [1998], Goswami et al. [2002])<sup>2</sup>, while others describe currency and financial crises as nonlinear processes (*e.g.* Jeanne and Masson [2000], Chang and Velasco [2001]). In growth economics, a number of theoretical models suggest that economic growth is a nonlinear process with the economy bouncing back and forth between different regimes (*e.g.* Zilibotti [1995], Peretto [1999], Matsuyama [1999], Galor and Weil [2000])<sup>3</sup>. Theoretical models in finance highlight heterogeneous expectations (*e.g.* Brock and Hommes [1998], De Grauwe and Grimaldi [2005]), heterogeneity in investors’ objectives (*e.g.* Peters [1994]), and herd behaviour (*e.g.* Lux [1995]) as some of the sources of nonlinearity in asset prices.

If economic and financial variables exhibit nonlinear behaviour, the standard unit root tests that are based on a linear AR process will have low power. Two recent papers, Sollis et al. [2002] and Kapetanios et al. [2003], address this issue by developing formal unit root tests against the alternative of nonlinear mean reversion. Both papers examine the unit root hypothesis against the nonlinear STAR (Smooth Transition AutoRegressive) alternative and show that, under the null hypothesis, the distribution of the respective tests is not normal. As a result the two papers employ Monte Carlo simulations to obtain critical values. The main difference between the two tests is that Sollis et al. [2002] use a logistic transition function (LSTAR) while Kapetanios et al. [2003] use an exponential transition function (ESTAR).

---

<sup>1</sup>For a review of the various unit root tests see, for example, Breitung and Pesaran [2007] and Cerrato and Sarantis [2007b].

<sup>2</sup>For empirical studies on nonlinear exchange rate models, see Michael et al. [1997], Sarantis [1999], Taylor et al. [2001], Rapach and Wohar [2003] among others.

<sup>3</sup>A number of authors have also undertaken empirical investigations of nonlinear growth models; see, for example, Fiaschi and Lavezzi [2007], Liu and Stengos [1999], Durlauf and Johnson [1995].

However both these nonlinear unit root tests are univariate and, consequently, will still suffer from low power in the case of small samples. In this paper we extend the [Kapetanios et al. \[2003\]](#) nonlinear unit root test to a panel context in order to address the low power problem of univariate tests. Since heterogeneous cross section dependence tends to be important in most empirical applications, we employ the [Pesaran \[2007\]](#) panel unit root framework that enables us to account for heterogeneous cross section dependence in a novel way. [Pesaran \[2007\]](#) shows that the individual CADF (Cross Augmented Dickey Fuller) and the panel statistic (CIPS) have non-normal distributions, so their critical values (for different  $N$  and  $T$ ) are obtained by Monte Carlo simulations. The panel unit root test proposed by [Pesaran \[2007\]](#) differs from other tests such as [Choi \[2001\]](#) and [Hadri \[2000\]](#) in that the latter all assume that individual time series are independent, and thus, cross section dependence is not considered. [Pesaran \[2007\]](#) shows that cross sectional dependence can be accounted for by augmenting the standard DF regression with the cross section averages of lagged levels and first differences of the individual series.

4

In this paper we propose a novel nonlinear panel unit root test that extends both the univariate nonlinear tests and the linear panel unit root tests, thus filling an important gap in the existing literature. Our test also allows for cross section dependence, and can be computed using basic OLS linear regression, and thus does not require any programming.<sup>5</sup> Since the panel nonlinear statistic has a non-normal distribution, we use Monte Carlo simulations to analyse the size and power of the test under different scenarios, and we calculate critical values which can be used in future applications of the test. Theorems 1 and 2 show that the distribution of the proposed test statistics is free of nuisance parameters. We finally illustrate the applicability of our test by applying it to a panel of bilateral real exchange rates.

The rest of the paper is organised as follows. Section 2 specifies the nonlinear dynamic panel model with cross section dependence. Section 3 derives the individual and panel nonlinear unit root tests, and then uses stochastic simulations to obtain the distributions of these statistics and critical values. Section 4 analyses the size and power of the panel nonlinear unit root test under alternative scenarios and compares the results to the performance of the linear [Pesaran \[2007\]](#) test. Section 5 reports the results from an application to real exchange rates, while [section 6](#) concludes.

## 2 A Nonlinear Dynamic Panel with Cross Section Dependence

Suppose the observation  $y_{it}$  on the  $i^{\text{th}}$  cross section unit at time  $t$  is generated according to the dynamic nonlinear heterogeneous panel ESTAR model below:

$$y_{it} = \beta_i y_{i,t-1} + \nu_i y_{i,t-1} Z(\theta_i; y_{i,t-d}) + u_{it}, \quad t = 1, \dots, T, \quad i = 1, \dots, N, \quad (1a)$$

---

<sup>4</sup>Another way to account for cross section dependence can be found in [Ng \[2008\]](#). However, [Ng \[2008\]](#) simply estimates the proportion of the panel that has a unit root and does not provide any information about the whole panel, conversely to the other tests. In addition, it is not a nonlinear test.

<sup>5</sup>[Chiang et al. \[2007\]](#) also proposes a nonlinear panel unit root test, but this test is based on an extension of the IPS test ([Im et al. \[2003\]](#)) and, therefore, does not account for cross section dependence which is so crucial in empirical applications.

where initial value,  $y_{i0}$ , is given, and the error term,  $u_{it}$ , has the one-factor structure:

$$u_{it} = \gamma_i f_t + \varepsilon_{it}, \quad (1b)$$

$$(\varepsilon_{it})_t \sim i.i.d.(0, \sigma_i^2), \quad (1c)$$

in which  $f_t$  is the unobserved common effect, and  $\varepsilon_{it}$  is the individual-specific (idiosyncratic) error. Following the literature on STAR models, the transition function adopted here is of the exponential form, *i.e.*,

$$Z(\theta_i; y_{i,t-d}) = 1 - \exp(-\theta_i y_{i,t-d}^2), \quad (1d)$$

where we assume that  $\theta_i \geq 0$ , and  $d \geq 1$  is the delay parameter. To begin with we assume that  $y_{it}$  is a mean zero stochastic process. We discuss processes with non-zero mean later. To simplify the model and following the existing literature, the delay parameter  $d$  is set to be equal to one and [Equation 1a](#)–[Equation 1d](#) are rewritten in first difference form as:

$$\Delta y_{it} = \phi_i y_{i,t-1} + \nu_i y_{i,t-1} [1 - \exp(-\theta_i y_{i,t-1}^2)] + \gamma_i f_t + \varepsilon_{it}, \quad (2)$$

where  $\phi_i = -(1 - \beta_i)$ . Assuming  $\phi_i = 0$ <sup>6</sup>, [Equation 2](#) can be rewritten as:

$$\Delta y_{it} = \nu_i y_{i,t-1} [1 - \exp(-\theta_i y_{i,t-1}^2)] + \gamma_i f_t + \varepsilon_{it}. \quad (3)$$

Using [Equation 3](#), we are interested in testing the hypothesis:

$$H_0 : \theta_i = 0 \forall i, \quad (4)$$

against the possibly heterogeneous alternatives:

$$H_1 : \begin{cases} \theta_i > 0 & \text{for } i = 1, \dots, N_1, \\ \theta_i = 0 & \text{for } i = N_1 + 1, \dots, N. \end{cases}$$

**Remark 1:** The alternative hypothesis above implies that some units are generated by a stationary ESTAR model but it also allows a proportion of units being a unit root process.

The following assumptions are introduced:

**Assumption 1:**  $\frac{N_1}{N} \rightarrow q$  as  $N \rightarrow \infty$ , with  $0 < q \leq 1$  under the alternative hypothesis [6](#)<sup>7</sup>.

**Assumption 2:**  $(\varepsilon_{it})$  are independently distributed for all  $i = 1, \dots, N$  and  $t = 1, \dots, T$ , with zero mean, constant variance  $\sigma_i^2$ , and finite fourth order moment.

**Assumption 3:**  $f_t$  is serially uncorrelated with zero mean, constant variance  $\sigma_f^2$ , and finite fourth moment. (Without loss of generality  $\sigma_f^2$  will be set equal to unity.)

**Assumption 4:**  $\varepsilon_{it}$ ,  $f_t$ , and  $\gamma_i$  are independently distributed for all  $i$ .

---

<sup>6</sup>It follows the practice in the literature (*e.g.* [Balke and Fomby \[1997\]](#), in the context of TAR models and [Michael et al. \[1997\]](#) in the context of ESTAR models).

<sup>7</sup>As noted in [Im et al. \[2003\]](#) this condition is necessary for the consistency of the panel unit root tests.

**Assumption 5:** Following Pesaran [2006], we define the weights  $(\omega_i)$  having the following properties:

$$\omega_N = O\left(\frac{1}{N}\right); \quad \sum_{i=1}^N \omega_i = 1; \quad \sum_{i=1}^N |\omega_i| < K < \infty; \quad \sum_{i=1}^N \omega_i^2 = O\left(\frac{1}{N}\right).$$

**Assumption 6:** Let  $\bar{\gamma} = \frac{1}{N} \sum_{i=1}^N \gamma_i$ . We assume  $\bar{\gamma} \neq 0$  for a fixed  $N$ .

**Assumption 7:** Following Pesaran and Tosetti [2007] we consider

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \gamma_i^2 = 0.$$

### 3 Nonlinear Unit Root Tests with Serially Uncorrelated Errors

Assumptions 1 and 2 together imply that the composite error,  $u_{it}$ , is serially uncorrelated. This restriction will be relaxed in subsection 3.3. Assumption 7 considers the degree of cross sectional dependence in our panel and it is useful for the development of the asymptotics. We shall relax it in our Monte Carlo experiments.<sup>8</sup>

#### 3.1 Individual NCADF Test

Testing the null hypothesis 4 directly is not feasible, since  $\nu_i$  is not identified under the null<sup>9</sup>. To overcome this problem, we follow Luukkonen et al. [1988], and derive a  $t$  type test statistic. Using Taylor expansion on Equation 3, under the null hypothesis, the following auxiliary regression is obtained:

$$\Delta y_{it} = a_i + b_i y_{i,t-1}^3 + \gamma_i f_t + e_{it}. \quad (5)$$

**Lemma 1:** If Assumptions 2–7 are satisfied, using the same approach as in Pesaran [2006], then the common factor  $f_t$  can be approximated by:

$$f_t \approx \frac{1}{\bar{\gamma}_\omega} \Delta \bar{y}_{\omega,t} - \frac{b}{\bar{\gamma}_\omega} \overline{y_{\omega,t-1}^3}, \quad (6)$$

where  $\Delta \bar{y}_{\omega,t} = \sum_{i=1}^N \omega_i \Delta y_{i,t}$ ,  $\overline{y_{\omega,t-1}^3} = \sum_{i=1}^N \omega_i y_{i,t-1}^3$ ,  $\bar{\gamma}_\omega = \sum_{i=1}^N \omega_i \gamma_i$ , and  $b$  is the mean of  $b_i$ .

**Proof:** see Appendix A.1.

Therefore, it follows that Equation 5 can be written as the following nonlinear cross sectionally augmented DF (NCADF) regression:

$$\Delta y_{it} = a_i + b_i y_{i,t-1}^3 + c_i \Delta \bar{y}_t + d_i \bar{y}_{t-1}^3 + e_{it} \quad (7)$$

<sup>8</sup>We wish to thank Ron Smith for calling our attention on this restriction.

<sup>9</sup>See for example Davies [1987].

The idea is, given the framework above, to develop a unit root test in the heterogeneous panel model based on [Equation 7](#). Extending the idea in [Kapetanios et al. \[2003\]](#), we suggest using [Equation 7](#) and the  $t$ -statistic on  $b_i$ , that is denoted by:

$$t_i(N, T) = \frac{\hat{b}_i}{s.e.(\hat{b}_i)}, \quad (8)$$

where  $\hat{b}_i$  is the OLS estimate of  $b_i$ , and  $s.e.(\hat{b}_i)$  its associated standard error. Denote the student statistic on the ratio of  $b_i$  in [Equation 7](#) as:

$$t_i(N, T) = \frac{y_{i,-1}^3{}' M \Delta y_i}{(\Delta y_i' M \Delta y_i)^{\frac{1}{2}} (y_{i,-1}^3{}' M y_{i,-1}^3)^{\frac{1}{2}}}, \quad (9)$$

where  $\Delta y_i = (\Delta y_{i1}, \Delta y_{i2}, \dots, \Delta y_{iT})'$ ,  $y_{i,-1}^3 = (y_{i,0}^3, y_{i,1}^3, \dots, y_{i,T-1}^3)'$ ,  $M$  the projection matrix onto  $\delta(X)$ , the orthogonal complement of the span of  $X$ ,  $M = I_T - X(X'X)^{-1}X'$ ,  $X = (\tau, \Delta \bar{y}, \bar{y}_{-1}^3)$ ,  $\tau = (1, 1, \dots, 1)'$ ,  $\Delta \bar{y} = (\Delta \bar{y}_1, \Delta \bar{y}_2, \dots, \Delta \bar{y}_T)'$ ,  $\bar{y}_{-1}^3 = (\bar{y}_0^3, \bar{y}_1^3, \dots, \bar{y}_{T-1}^3)$ . The critical values of the NCADF test can be computed by stochastic simulation for any fixed  $T > 3$ , and for given distributional assumptions for the random variables  $(\varepsilon, f)$ .<sup>10</sup>

It should be noted that, as in [Pesaran \[2007\]](#), the exact null distribution of the individual test statistic  $t_i(N, T)$  defined in [Equation 9](#) is affected by the nuisance parameters. However, this distribution depends on the nuisance parameters only through their effects on the matrix  $M$ . The following theorem shows that this dependence vanishes as  $N \rightarrow \infty$ .

**Theorem 1** *Suppose the cross-section mean of the initial observations  $\bar{y}_0$  is set to zero. Then,*

$$\bar{t}(N, T) = \frac{1}{N} \sum_{i=1}^N t_i(N, T) = \frac{1}{N} \sum_{i=1}^N \tau_i(N, T) + R,$$

where, under Assumptions 1–7, as  $N \rightarrow \infty$  and  $T$  is fixed,  $\tau_i(N, T)$  converges to a distribution which is free of nuisance parameters, and where  $R$  converges to zero.

See proof in [Appendix A.2](#).

**Theorem 2** *Suppose  $\bar{y}_0 = 0$ , and that Assumptions 1–7 hold. Then, as  $(N, T) \rightarrow \infty$  simultaneously such that  $N/T$  tends to a finite constant,  $\tau_i(N, T)$  converges in distribution to*

$$\frac{\int W_i^3 dW_i - q' \Psi^{-1} h}{\sqrt{\int W_i^6 - h' \Psi^{-1} h}},$$

<sup>10</sup>To accommodate stochastic processes with non-zero means we follow [Kapetanios et al. \[2003\]](#). In the case when the process has non-zero mean, we use demeaned and detrended data, i.e. when  $x_t = \mu + \sigma t + y_t$ , we use  $y_t = x_t - \hat{\mu} + \hat{\sigma} t$ , where  $\hat{\mu}$  and  $\hat{\sigma}$  are the OLS estimators of  $\mu$  and  $\sigma$  (see [Kapetanios et al. \[2003\]](#) for further details).

where

$$\Psi = \begin{pmatrix} 1 & \int W_f^3 \\ \int W_f^3 & \int W_f^6 \end{pmatrix},$$

$$q = \begin{pmatrix} W_i(1) \\ \int W_f^3 dW_i \end{pmatrix}, \quad h = \begin{pmatrix} \int W_i^3 \\ \int W_f^3 W_i^3 \end{pmatrix},$$

using the short-hand notation

$$\int W_i^3 dW_i = \int_0^1 W_i(t)^3 dW_i(t),$$

$$\int W_i^6 = \int_0^1 W_i(t)^6 dt,$$

etcetera, where  $W_i$  and  $W_f$  are standard Wiener processes that are independent of each other.

See proof in Appendix A.3.

Figure 1 displays the simulated cumulative distribution function of the individual NCADF statistic under the null hypothesis using 50,000 replications for  $N = 100$  and  $T = 500$ . For comparison the simulated cumulative distribution function of Pesaran CADF statistic is also provided. The series  $y_{it} = y_{i,t-1} + f_t + u_{it}$ , for  $i = 1, 2, \dots, 100$ , and  $t = -50, -49, \dots, 1, 2, \dots, 500$  were first generated from  $y_{i,-50} = 0$ , with  $f_t$  and  $u_{it}$  as i.i.d.  $N(0,1)$ . Then 50,000 NCADF regressions of  $\Delta y_{it}$  on  $y_{i,t-1}^3$ ,  $\Delta \bar{y}_t$ , and  $\bar{y}_{t-1}^3$ .  $\Delta \bar{y}_t$ , and  $\bar{y}_{t-1}^3$  were computed over the sample  $t = 1, 2, \dots, 500$ . Figure 1 plots the ordered values of the OLS  $t$ -ratios of  $y_{i,t-1}^3$  in these regressions.

Not surprisingly the nonlinear CADF distribution, as the Pesaran's CADF distribution, is more skewed to the left as compared to the standard DF distribution. This is clearly reflected in the critical values of the distributions summarized in Table 1. Critical values of the individual nonlinear CADF distribution for values of  $T$  and  $N$  in the range of 10 to 200 are given Table 11 in Appendix B.1.

Table 1: Critical Values of the DF, Pesaran's CADF, and nonlinear CADF Distributions ( $N = 100, T = 500, 50,000$  replications)

	1%	2.5%	5%	10%
DF	-2.60	-2.23	-1.94	-1.61
Pesaran's CADF	-3.80	-3.49	-3.22	-2.91
Nonlinear CADF	-3.72	-3.41	-3.15	-2.85

The nonlinear CADF distribution, like the Pesaran's CADF distribution and the standard DF distribution, departs from normality in two important respects: it has a substantially negative mean and its standard deviation is less than unity, although not by a large amount. The simulated density functions of the standardized NCADF, computed with  $N = 100, T = 500$ , and 50,000 replications are displayed in Figure 2. The mean, standard deviation, skewness and Kurtosis -3 coefficients of the NCADF and the Pesaran's CADF distributions are reported in Table 2. They are quite small, although statistically highly significant.

Figure 1: Cumulative Distribution Function of Pesaran's Cross Sectionally Augmented DF and Nonlinear Cross Sectionally Augmented DF Statistics




Figure 2: Simulated Density Function of the Standardized NCADF<sub>i</sub> and the Standardized Pesaran's CADF<sub>i</sub> Distributions as Compared to the Normal Density






Table 2: Moments of the CADF distributions

	Pesaran's CADF	NCADF
Mean	-1.80	-1.83
Standard deviation	0.90	0.83
Skewness	0.20	0.28
Kurtosis -3	0.19	0.77

Table 3: Moments of the CIPS distributions

	Pesaran's CIPS	NCIPS
Mean	-1.80	-1.83
Standard deviation	0.17	0.12
Skewness	-0.10	-0.068
Kurtosis -3	-1.67	-1.45

Since cross sectional dependence in panel data is widely known now to be a serious problem, in the next sections we shall be using Equation 7 to develop a unit root test to test for the null hypothesis of unit root against an ESTAR stationary alternative.

### 3.2 Panel Nonlinear CADF Test

Following Pesaran [2007], we suggest using the  $t$ -statistic in Equation 9 to construct a panel unit root test by averaging the individual test statistics:

$$\bar{t}(N, T) = \frac{1}{N} \sum_{i=1}^N t_i(N, T) \quad (10)$$

This is a nonlinear cross sectionally augmented version of the IPS test based (NCIPS). The test statistic defined in Equation 10 can also be extended to the case where serial correlation is present in the data. In this particular case, one may include, in the model, lags of the left hand side variable after using an information criteria to select the lag order.


We simulated the distribution of NCIPS setting  $N = 100$ ,  $T = 500$ , and using 50,000 replications. The simulated density functions of the NCIPS and the Pesaran's CIPS Statistics are displayed in Figure 3. Both the densities show marked departures from normality. The skewness and Kurtosis -3 coefficients of the NCIPS and the Pesaran's CIPS distributions are reported in Table 3. The critical values of the nonlinear CIPS test are given in Table 12 in Appendix B.2.

### 3.3 The Serially Correlated Errors Case

Serial correlation can be incorporated in the model in a variety of different ways. In what follows, we use the model in Equation 5 and specify the serial correlation structure as:

$$u_{it} = \rho_i u_{i,t-1} + \eta_{it}. \quad (11)$$

Figure 3: Simulated Density Function of the NCIPS Statistic and the Pesaran's CIPS Distributions



We first model serial correlation as above and thereafter cross section dependence as:

$$\eta_{it} = \gamma_i f_t + \varepsilon_{it}. \quad (12)$$

Using Equation 5 jointly with Equation 11 above we obtain:

$$\Delta y_{it} = a_i(1 - \rho_i) + b_i(1 - \rho_i)y_{i,t-1}^3 + \rho_i \Delta y_{i,t-1} + b_i \rho_i \Delta (y_{i,t-1}^3) + \eta_{it}. \quad (13)$$

And substituting Equation 12 into Equation 13

$$\Delta y_{it} = a_i(1 - \rho_i) + b_i(1 - \rho_i)y_{i,t-1}^3 + \rho_i \Delta y_{i,t-1} + b_i \rho_i \Delta (y_{i,t-1}^3) + \gamma_i f_t + \varepsilon_{it}. \quad (14)$$

Using Equation 14 and the same approach as in Appendix A.1, one can obtain a proxy for  $f_t$  using the following set of variables:

$$\left\{ \overline{y_{t-1}^3}, \overline{\Delta y_{t-1}^3} \right\}$$

By generalizing this to an AR(p) error terms framework, we suggest using the following general nonlinear CADF regression:

$$\Delta y_{it} = a_i + b_{i0} y_{i,t-1}^3 + d_{i0} \overline{y_{t-1}^3} + \sum_{j=1}^p d_{ij} \overline{\Delta y_{t-j}^3} + \sum_{j=1}^p \delta_{ij} \Delta y_{i,t-j} + e_{it}. \quad (15)$$

Information criteria can be used to choose the length of  $p$ .

## 4 Small Sample Analysis

In this section we assess the size and power of the nonlinear panel test defined in Equation 10 under different scenarios. About the power of the test, we firstly look at it in the

case of weak and strong cross sectional dependence, but no autocorrelation structure for the error term. In the next section, we generalise this scenario by allowing an autocorrelation specification for the error term and weak-strong cross sectional dependence. For comparison, in all the experiments we also report the size and power of Pesaran [2007] test when a nonlinear DGP is considered.

The data generating process (DGP) considered is the following Panel ESTAR:

$$\Delta y_{it} = \nu_i y_{i,t-1} [1 - \exp(-\theta_i y_{i,t-1}^2)] + \gamma_i f_t + \varepsilon_{it}, \quad (16a)$$

$$f_t \sim i.i.d.N(0, 1), \quad (16b)$$

$$\varepsilon_{it} \sim i.i.d.N(0, \sigma_i^2), \quad (16c)$$

$$\sigma_i^2 \sim i.i.d.U[0.5, 1.5]. \quad (16d)$$

with  $i = 1, 2, \dots, N$  and  $t = -51, -50, \dots, 1, 2, \dots, T$ . We fix  $\nu_i = -1$  for all  $i$ . The choice of the cross sectional dependence parameters  $\gamma_i$  depends on whether we wish to impose Assumption 7 or not.

This DGP is considered in two cases for Assumption 7. The first under Assumption 7 and therefore follows our theoretical analysis in Appendix A. We consider two scenarios for cross sectional dependence, namely weak cross sectional dependence  $\gamma_i \sim i.i.d.U[0, 0.20] \times 30/i$ , and strong cross sectional dependence  $\gamma_i \sim i.i.d.U[1, 3] \times 30/i$ .

The second DGP relaxes Assumption 7. We also consider two scenarios for cross sectional dependence, namely weak cross sectional dependence  $\gamma_i \sim i.i.d.U[0, 0.20]$ , and strong cross sectional dependence  $\gamma_i \sim i.i.d.U[1, 3]$ .

## 4.1 Size Distortion Analysis

In our size analysis below, we generate data by setting  $\theta = 0$  for all  $i$ . Size is computed at the 5% nominal significance level. The number of replications is set to 5,000. The standard error of the computed size is 0.0031.

Under assumption 7, results for the size are reported in Table 4. Results for the size when Assumption 7 is relaxed are presented in Appendix C, Table 13.

As expected the test statistics seems to be correctly sized in the case of weak cross section dependence and as  $N$  and  $T$  become large.<sup>11</sup> Let us now consider the size of the test when Assumption 7 is relaxed. The test seems to have an acceptable size for large cross section dimension and somehow slightly undersized with respect to Pesaran [2007] test.

## 4.2 Power Analysis

In this section we assess the power of the test defined in Equation 10 under the same DGP as above but we consider the cases of weak and strong alternatives, namely we assume for the weak alternative:

$$\theta_i = \begin{cases} 0 & \text{for } i = 1, \dots, N/2, \\ 0.01 & \text{for } i = N/2 + 1, \dots, N, \end{cases} \quad (17a)$$

while for the strong alternative:

$$\theta_i = \begin{cases} 0 & \text{for } i = 1, \dots, N/2, \\ 0.05 & \text{for } i = N/2 + 1, \dots, N. \end{cases} \quad (17b)$$

---

<sup>11</sup>Note that Assumption 7 implies weak cross section dependence.

The power is computed at the 5% nominal significance level. Under assumption 7, results are reported in [Table 5](#) (weak alternative) and [Table 6](#) (strong alternative). Results for the power when Assumption 7 is relaxed are presented in [Appendix C](#), [Table 14](#) (weak alternative) and [Table 15](#) (strong alternative).

The test we propose seems to have stronger power than the [Pesaran \[2007\]](#) test when the true DGP is nonlinear regardless of Assumption 7.

### 4.3 Serial Correlated Errors Case

In this section we analyze size and power of the proposed test when serial correlation is incorporated into the DGP. We consider positive serial correlation. The error terms ( $\varepsilon_{it}$ ) were generated as:

$$\varepsilon_{it} = \rho_i \varepsilon_{i,t-1} + \zeta_{it}, \quad (18a)$$

$$\zeta_{it} \sim i.i.d.N(0, \sigma_i^2), \quad (18b)$$

$$\sigma_i^2 \sim i.i.d.U[0.5; 1.5], \quad (18c)$$

$$\rho_i \sim i.i.d.U[0.2; 0.4] \quad \text{in the case of positive correlation,} \quad (18d)$$

$$\rho_i \sim i.i.d.U[-0.4; -0.2] \quad \text{in the case of negative correlation.} \quad (18e)$$

We only consider here the power analysis for the case of strong alternative:

$$\theta_i = \begin{cases} 0 & \text{for } i = 1, \dots, N/2, \\ 0.05 & \text{for } i = N/2 + 1, \dots, N. \end{cases}$$

and strong cross sectional dependence  $\gamma_i \sim i.i.d.U[1, 3]$ . The size and power are computed at 5% nominal significance level and it are based on the following nonlinear CADF regression:

$$\Delta y_{it} = a_i + b_i y_{i,t-1}^3 + d_{i,0} \overline{y_{t-1}^3} + d_{i,1} \overline{\Delta y_{t-1}^3} + \delta_{i,1} \Delta y_{i,t-1} + e_{it}, \quad (19)$$

$$i = 1, 2, \dots, N; t = 1, 2, \dots, T; \bar{y}_t = \frac{1}{N} \sum_{i=1}^N y_{it}.$$

The test is computed as:

$$\bar{t}(N, T) = \frac{1}{N} \sum_{i=1}^N t_{iNL}(N, T) \quad (20)$$

where  $\bar{t}(N, T)$  is the OLS  $t$ -ratio of  $b_i$  in the above nonlinear ADF regression. The number of simulations is set equal to 50,000. [Table 7](#) shows the results for the size of the tests when Assumption 7 is respected. [Table 16](#) in [Appendix C](#) shows the results for the size of the tests when Assumption 7 is relaxed. Both tests have a good size with the Pesaran's test being consistently oversized.

In [Table 8](#) we show results on the power of the test in the case when positive as well as negative serial correlation is present in the DGP, under Assumption 7. [Table 17](#) in [Appendix C](#) presents the same results when Assumption 7 is relaxed. For panels of a moderate size, the gain in power from using the nonlinear panel unit root test with respect to the Pesaran's test is evident regardless of Assumption 7.

Table 4: Size: Case of no serial correlation

Sizes		Weak Cross Section Dependence				Strong Cross Section Dependence			
$N$	$T$	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS
10	10	0.0482	0.0448	0.0554	0.0478	0.0456	0.0434	0.0594	0.0524
10	20	0.0502	0.0432	0.0488	0.0420	0.0470	0.0474	0.0650	0.0480
10	30	0.0544	0.0490	0.0544	0.0418	0.0504	0.0560	0.0622	0.0488
10	50	0.0562	0.0446	0.0476	0.0446	0.0446	0.0456	0.0630	0.0520
10	100	0.0456	0.0486	0.0492	0.0428	0.0486	0.0458	0.0670	0.0506
20	10	0.0502	0.0538	0.0476	0.0446	0.0468	0.0476	0.0544	0.0482
20	20	0.0462	0.0496	0.0504	0.0398	0.0428	0.0406	0.0558	0.0494
20	30	0.0542	0.0498	0.0548	0.0432	0.0504	0.0470	0.0592	0.0456
20	50	0.0480	0.0556	0.0524	0.0446	0.0474	0.0446	0.0612	0.0418
20	100	0.0448	0.0470	0.0572	0.0464	0.0510	0.0482	0.0554	0.0394
30	10	0.0440	0.0482	0.0512	0.0444	0.0554	0.0514	0.0568	0.0448
30	20	0.0524	0.0478	0.0486	0.0446	0.0454	0.0448	0.0530	0.0420
30	30	0.0542	0.0534	0.0608	0.0454	0.0508	0.0476	0.0616	0.0368
30	50	0.0480	0.0484	0.0554	0.0412	0.0510	0.0476	0.0612	0.0426
30	100	0.0486	0.0456	0.0646	0.0462	0.0474	0.0466	0.0620	0.0384
50	10	0.0478	0.0548	0.0488	0.0482	0.0530	0.0490	0.0524	0.0414
50	20	0.0516	0.0502	0.0438	0.0394	0.0460	0.0476	0.0554	0.0414
50	30	0.0602	0.0494	0.0530	0.0460	0.0492	0.0444	0.0530	0.0406
50	50	0.0502	0.0536	0.0486	0.0422	0.0484	0.0462	0.0570	0.0386
50	100	0.0506	0.0512	0.0506	0.0466	0.0476	0.0534	0.0538	0.0364
100	10	0.0464	0.0468	0.0520	0.0490	0.0474	0.0452	0.0480	0.0456
100	20	0.0512	0.0440	0.0558	0.0456	0.0532	0.0494	0.0470	0.0420
100	30	0.0548	0.0500	0.0474	0.0454	0.0444	0.0432	0.0532	0.0424
100	50	0.0500	0.0508	0.0430	0.0444	0.0478	0.0480	0.0598	0.0396
100	100	0.0592	0.0556	0.0472	0.0512	0.0486	0.0516	0.0564	0.0352

Table 5: Power: Case of no serial correlation and weak alternative

Sizes		Weak Cross Section Dependence				Strong Cross Section Dependence			
$N$	$T$	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS
10	10	0.0430	0.0454	0.0592	0.0662	0.3100	0.1396	0.5042	0.5278
10	20	0.0556	0.0456	0.0770	0.0870	0.7326	0.3546	0.8980	0.8884
10	30	0.0586	0.0586	0.1158	0.1370	0.8102	0.5646	0.9424	0.9354
10	50	0.0702	0.0482	0.2324	0.2702	0.8354	0.7240	0.9554	0.9498
10	100	0.1096	0.0704	0.6450	0.7514	0.8716	0.8210	0.9662	0.9650
20	10	0.0540	0.0490	0.0672	0.0680	0.1446	0.1280	0.2784	0.3992
20	20	0.0476	0.0478	0.0994	0.1180	0.5956	0.3064	0.8014	0.8222
20	30	0.0502	0.0502	0.1510	0.1868	0.7472	0.4798	0.9384	0.9528
20	50	0.0602	0.0514	0.2558	0.3584	0.8394	0.6780	0.9812	0.9844
20	100	0.0958	0.0678	0.7548	0.8972	0.8840	0.8286	0.9870	0.9872
30	10	0.0502	0.0490	0.0708	0.0734	0.0808	0.0932	0.1480	0.2496
30	20	0.0472	0.0398	0.0938	0.1348	0.3962	0.2250	0.6066	0.6760
30	30	0.0542	0.0434	0.1486	0.2128	0.6366	0.3870	0.8640	0.8960
30	50	0.0530	0.0508	0.3236	0.5038	0.7846	0.5968	0.9798	0.9860
30	100	0.0718	0.0504	0.8318	0.9704	0.8742	0.7802	0.9930	0.9938
50	10	0.0468	0.0450	0.0700	0.0808	0.0582	0.0602	0.0864	0.1228
50	20	0.0490	0.0484	0.1098	0.1700	0.1968	0.1268	0.3360	0.3974
50	30	0.0536	0.0472	0.1842	0.3010	0.4092	0.2448	0.6444	0.7162
50	50	0.0616	0.0528	0.3594	0.6578	0.6456	0.4162	0.9216	0.9642
50	100	0.0742	0.0632	0.9204	0.9970	0.8138	0.6646	0.9990	0.9992
100	10	0.0502	0.0428	0.0606	0.0838	0.0454	0.0512	0.0568	0.0782
100	20	0.0484	0.0446	0.1156	0.2338	0.0780	0.0662	0.1596	0.2064
100	30	0.0550	0.0518	0.1970	0.4484	0.1734	0.1174	0.3314	0.4312
100	50	0.0544	0.0534	0.4418	0.8510	0.3486	0.1864	0.7108	0.8500
100	100	0.0642	0.0534	0.9734	1.000	0.6278	0.3768	0.9990	0.9998

Table 6: Power: Case of no serial correlation and strong alternative

Sizes		Weak Cross Section Dependence				Strong Cross Section Dependence			
$N$	$T$	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS
10	10	0.0548	0.0464	0.0818	0.0958	0.5288	0.1278	0.7820	0.6412
10	20	0.0640	0.0514	0.1904	0.2632	0.8148	0.3736	0.9510	0.9176
10	30	0.0948	0.0660	0.3760	0.4736	0.8492	0.5960	0.9608	0.9434
10	50	0.1410	0.0778	0.7734	0.8674	0.8730	0.7624	0.9628	0.9544
10	100	0.2028	0.1152	0.9968	0.9994	0.8768	0.8330	0.9632	0.9604
20	10	0.0458	0.0466	0.0896	0.1136	0.3960	0.1382	0.6510	0.6352
20	20	0.0578	0.0506	0.2270	0.3548	0.7926	0.3612	0.9722	0.9582
20	30	0.0778	0.0558	0.4306	0.6358	0.8542	0.5506	0.9832	0.9772
20	50	0.0912	0.0646	0.8684	0.9668	0.8860	0.7716	0.9870	0.9858
20	100	0.1312	0.0766	1.000	1.000	0.8950	0.8484	0.9882	0.9886
30	10	0.0572	0.0542	0.1018	0.1292	0.2792	0.1252	0.4846	0.5494
30	20	0.0552	0.0472	0.2492	0.4422	0.7350	0.3070	0.9576	0.9612
30	30	0.0606	0.0528	0.4874	0.7946	0.8402	0.5186	0.9916	0.9910
30	50	0.0788	0.0584	0.9250	0.9956	0.8764	0.7326	0.9946	0.9948
30	100	0.1276	0.0774	1.000	1.000	0.8986	0.8426	0.9966	0.9976
50	10	0.0504	0.0490	0.0998	0.1626	0.1428	0.0970	0.2610	0.3744
50	20	0.0576	0.0438	0.2712	0.5928	0.5546	0.2518	0.8492	0.9088
50	30	0.0640	0.0508	0.5684	0.9134	0.7440	0.4104	0.9872	0.9944
50	50	0.0760	0.0624	0.9714	1.000	0.8380	0.6052	0.9980	0.9988
50	100	0.1102	0.0742	1.000	1.000	0.8830	0.7830	0.9992	0.9994
100	10	0.0534	0.0502	0.1130	0.2044	0.0728	0.0632	0.1334	0.2124
100	20	0.0524	0.0538	0.3248	0.7712	0.2772	0.1256	0.5744	0.7972
100	30	0.0616	0.0538	0.6592	0.9916	0.4774	0.2256	0.9172	0.9912
100	50	0.0680	0.0614	0.9962	1.000	0.6750	0.3754	1.000	1.000
100	100	0.0962	0.0668	1.000	1.000	0.8050	0.6154	1.000	1.000

Table 7: Size: Case of serial correlation

Sizes		Weak Cross Section Dependence				Strong Cross Section Dependence			
$N$	$T$	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS
10	10	0.0482	0.0448	0.0554	0.0378	0.0456	0.0434	0.0594	0.0524
10	20	0.0502	0.0432	0.0488	0.0420	0.0470	0.0474	0.0650	0.0480
10	30	0.0544	0.0490	0.0544	0.0418	0.0504	0.0560	0.0622	0.0488
10	50	0.0562	0.0446	0.0476	0.0446	0.0446	0.0456	0.0630	0.0520
10	100	0.0456	0.0486	0.0492	0.0428	0.0486	0.0458	0.0670	0.0506
20	10	0.0502	0.0538	0.0476	0.0446	0.0468	0.0476	0.0544	0.0482
20	20	0.0462	0.0496	0.0504	0.0398	0.0428	0.0406	0.0558	0.0494
20	30	0.0542	0.0498	0.0548	0.0432	0.0504	0.0470	0.0592	0.0456
20	50	0.0480	0.0556	0.0524	0.0446	0.0474	0.0446	0.0612	0.0418
20	100	0.0448	0.0470	0.0572	0.0464	0.0510	0.0482	0.0554	0.0394
30	10	0.0440	0.0482	0.0512	0.0444	0.0554	0.0514	0.0568	0.0448
30	20	0.0524	0.0478	0.0486	0.0446	0.0454	0.0448	0.0530	0.0420
30	30	0.0542	0.0534	0.0608	0.0454	0.0508	0.0476	0.0616	0.0368
30	50	0.0480	0.0484	0.0554	0.0412	0.0510	0.0476	0.0612	0.0426
30	100	0.0486	0.0456	0.0646	0.0462	0.0474	0.0466	0.0620	0.0384
50	10	0.0478	0.0548	0.0488	0.0482	0.0530	0.0490	0.0524	0.0414
50	20	0.0516	0.0502	0.0438	0.0394	0.0460	0.0476	0.0554	0.0414
50	30	0.0602	0.0494	0.0530	0.0460	0.0492	0.0444	0.0530	0.0406
50	50	0.0502	0.0536	0.0486	0.0422	0.0484	0.0462	0.0570	0.0386
50	100	0.0506	0.0512	0.0506	0.0466	0.0476	0.0534	0.0538	0.0364
100	10	0.0464	0.0468	0.0520	0.0490	0.0474	0.0452	0.048	0.0456
100	20	0.0512	0.0440	0.0558	0.0456	0.0532	0.0494	0.047	0.0420
100	30	0.0548	0.0500	0.0474	0.0454	0.0444	0.0432	0.0532	0.0424
100	50	0.0500	0.0508	0.0430	0.0444	0.0478	0.0480	0.0598	0.0396
100	100	0.0592	0.0556	0.0472	0.0512	0.0486	0.0516	0.0564	0.0352

Table 8: Power: Case of serial correlation and strong alternative

Sizes		Weak Cross Section Dependence				Strong Cross Section Dependence			
$N$	$T$	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS	$CIPS_i$	$NCIPS_i$	CIPS	NCIPS
10	10	0.0548	0.0464	0.0818	0.0958	0.5288	0.1278	0.7820	0.6412
10	20	0.0640	0.0514	0.1904	0.2632	0.8148	0.3736	0.9510	0.9176
10	30	0.0948	0.0660	0.3760	0.4736	0.8492	0.5960	0.9608	0.9434
10	50	0.1410	0.0778	0.7734	0.8674	0.8730	0.7624	0.9628	0.9544
10	100	0.2028	0.1152	0.9968	0.9994	0.8768	0.8330	0.9632	0.9604
20	10	0.0458	0.0466	0.0896	0.1136	0.3960	0.1382	0.6510	0.6352
20	20	0.0578	0.0506	0.2270	0.3548	0.7926	0.3612	0.9722	0.9582
20	30	0.0778	0.0558	0.4306	0.6358	0.8542	0.5506	0.9832	0.9772
20	50	0.0912	0.0646	0.8684	0.9668	0.8860	0.7716	0.9870	0.9858
20	100	0.1312	0.0766	1.000	1.000	0.8950	0.8484	0.9882	0.9886
30	10	0.0572	0.0542	0.1018	0.1292	0.2792	0.1252	0.4846	0.5494
30	20	0.0552	0.0472	0.2492	0.4422	0.7350	0.3070	0.9576	0.9612
30	30	0.0606	0.0528	0.4874	0.7946	0.8402	0.5186	0.9916	0.9910
30	50	0.0788	0.0584	0.9250	0.9956	0.8764	0.7326	0.9946	0.9948
30	100	0.1276	0.0774	1.000	1.000	0.8986	0.8426	0.9966	0.9976
50	10	0.0504	0.0490	0.0998	0.1626	0.1428	0.0970	0.2610	0.3744
50	20	0.0576	0.0438	0.2712	0.5928	0.5546	0.2518	0.8492	0.9088
50	30	0.0640	0.0508	0.5684	0.9134	0.7440	0.4104	0.9872	0.9944
50	50	0.0760	0.0624	0.9714	1.000	0.8380	0.6052	0.9980	0.9988
50	100	0.1102	0.0742	1.000	1.000	0.8830	0.7830	0.9992	0.9994
100	10	0.0534	0.0502	0.1130	0.2044	0.0728	0.0632	0.1334	0.2124
100	20	0.0524	0.0538	0.3248	0.7712	0.2772	0.1256	0.5744	0.7972
100	30	0.0616	0.0538	0.6592	0.9916	0.4774	0.2256	0.9172	0.9912
100	50	0.0680	0.0614	0.9962	1.000	0.6750	0.3754	1.000	1.000
100	100	0.0962	0.0668	1.000	1.000	0.8050	0.6154	1.000	1.000

## 5 An Empirical Application: Real Exchange Rates

In this section we apply our test to real exchange rates against the US dollar for twenty OECD countries over the period 1973Q1–1998Q2. The data set is the same used by [Murray and Papell \[2002, 2005\]](#).

Since the long run *Purchasing Power Parity* (PPP) relationship is one of the main components of theoretical international macroeconomic models, a large number of studies have tested this relationship by applying unit root tests to real exchange rates. Most of these studies show evidence of unit root behaviour in real exchange rates, which has become a puzzle in international finance. The growing literature on nonlinear exchange rates argues that transaction costs and frictions in financial markets may lead to nonlinear convergence in real exchange rates. Consequently, the non-mean reversion reported by linear unit root tests may be due to the fact that these tests are based on a mis-specified stochastic process.

The individual statistics for our unit root test are shown in [Table 9](#). For comparison purposes, we also report the statistics for the [Pesaran \[2007\]](#) test which accounts for cross section dependence but not for nonlinearity.

The [Pesaran \[2007\]](#) test rejects the unit root null hypothesis in only 1 out of 20 cases at all levels of significance. By contrast, the nonlinear test rejects the null in 2 cases at the 1% significance level, and in 5 cases at the 5% and 10% level. Hence our test rejects the unit root null more frequently and therefore yields stronger support for the long-run PPP.

As we argued above, univariate tests have low power and this problem is overcome by employing panel unit root tests. The results for our panel unit root test and the [Pesaran](#) panel unit root test are shown in [Table 10](#).

The contrast between the two panel statistics is rather strong. The [Pesaran \[2007\]](#)

Table 9: Individual Unit Root Tests for Real Dollar Exchange Rates

Country	Lag	NCADF	CADF
Australia	3	-2.1765	-1.6501
Austria	4	-2.2085	-2.1432
Belgium	4	-2.4220	-1.2380
Canada	6	-1.1528	-1.3575
Denmark	3	-3.3390**	-2.8699
Finland	7	-1.7015	-2.4148
France	4	-0.9386	-2.1170
Germany	4	-3.3166**	-2.6044
Greece	4	-0.1449	-2.1730
Ireland	6	-0.1855	-1.0970
Italy	4	-2.6717	-2.0218
Japan	3	-2.5943	-1.9477
Netherlands	4	-2.7076	-1.9930
N Zealand	3	-3.7296**	-3.8758***
Norway	7	-2.2595	-1.8869
Portugal	8	-1.9120	-0.6359
Spain	8	-1.6911	-2.1622
Sweden	8	-3.8830***	-1.5888
Switzerland	4	-5.1263***	-2.7768
UK	7	-2.5354	-2.0689
Critical Values ( $N = 20, T = 100$ ):			
1%		-3.74	-3.87
5%		-3.09	-3.24
10%		-2.80	-2.92

\*\* Statistics significant at 5% level.

\*\*\* Statistics significant at 1% level.

Rejection Rates of the Panel Unit Root Tests

Level	NCADF		CADF	
	H0	H1	H0	H1
1%	90%	10%	95%	5%
5%	75%	25%	95%	5%
10%	75%	25%	95%	5%

Table 10: Panel Unit Root Tests for Real Dollar Exchange Rates

Country	NCADF	CADF
Panel	-2.3348***	-2.0311
Critical Values ( $N = 20, T = 100$ ):		
1%	-2.24	-2.36
5%	-2.11	-2.20
10%	-2.03	-2.11

\*\*\* Statistic significant at 1% level.



test fails to reject the unit root null at all levels of significance, thus implying non-mean reversion. On the other hand, our nonlinear panel test rejects the unit root null for the panel of real exchange rates at all levels of significance, giving support to the long-run PPP. This evidence of nonlinear mean reversion in the OECD real exchange rates may suggest that previous evidence of non-mean reversion in real exchange rates might be due to using linear unit root tests.

## 6 Conclusion

A number of panel unit root tests allowing for cross section dependence have been proposed in the literature. In this paper we propose a nonlinear heterogeneous panel unit root test for testing the null hypothesis of unit-root processes against the alternative that allows a proportion of units to be generated by globally stationary ESTAR processes and a remaining non-zero proportion to be generated by unit root processes. The proposed test is simple to apply and accommodates both nonlinearity and cross sectional dependence. Our test is compared to the [Pesaran \[2007\]](#) linear test via Monte Carlo simulation exercises, and it is found that our test holds correct size and under the hypothesis that data are generated by globally stationary ESTAR processes has a better power than the Pesaran test. We also calculate critical values for varying cross section and time dimensions which can be used in future applications of our test.

We provide an application to a panel of bilateral real exchange rate series with the US dollar from the 20 major OECD countries. In contrast to the evidence obtained by linear tests, we find evidence of nonlinear mean-reversion in the real exchange rates for the whole OECD panel that gives support to the long run PPP hypothesis. Given the importance of the PPP in international macroeconomic models, our evidence suggests that the employment of nonlinear panel unit root tests may provide a solution to the PPP puzzle.

Given the growing literature of nonlinear models, we believe that the development of panel nonlinear unit root tests has large potential in macroeconomic and financial applications. Evidence indicates that different time series may follow different nonlinear specifications. Consequently, one could consider unit root tests with different types of transition functions that allow for asymmetric dynamic adjustment. Another extension would be to allow for different transition variables. Further applications of our tests and theoretical extensions are left for future work.

# Appendix

## A Proofs

### A.1 Lemma 1

We assume that the error term  $e_{it}$  in Equation 5 follows a stationary process, for all  $i$ , with summable auto-covariance given by:

$$e_{it} = \sum_{l=0}^{\infty} a_{il} \zeta_{i,t-l}, \quad (\text{A.1.1})$$

with  $\zeta_{i,t}$  being a zero mean random variable with variance matrix defined by  $I_{l+1}$  and finite fourth order moment. The variance of  $e_{it}$  is finite and given by:

$$\text{Var}(e_{it}) = \sum_{l=0}^{\infty} a_{il}^2 = \sigma_i^2 \leq \sigma^{*2} < \infty, \quad (\text{A.1.2})$$

for some constant  $\sigma^{*2}$ .

First note that following Pesaran [2006] and after using cross sectional averages to average across  $i$ , Equation 5 can be written as:

$$\Delta \bar{y}_{\omega,t} = \bar{a}_{\omega} + \overline{y_{\omega,t-1}^3} b + \bar{\gamma}_{\omega} f_t + \bar{e}_{\omega,t}, \quad (\text{A.1.3})$$

with  $\Delta \bar{y}_{\omega,t} = \sum_{i=1}^N \omega_i \Delta y_{i,t}$ ,  $\bar{a}_{\omega} = \sum_{i=1}^N \omega_i a_i$  and  $\bar{\gamma}_{\omega} = \sum_{i=1}^N \omega_i \gamma_i$ .

Assuming that  $\bar{\gamma}_{\omega} \neq 0$  for fixed  $T$ , then  $f_t$  can be approximated as follows:

$$f_t \approx \frac{1}{\bar{\gamma}_{\omega}} \Delta \bar{y}_{\omega,t} - \frac{1}{\bar{\gamma}_{\omega}} b \overline{y_{\omega,t-1}^3} - \frac{1}{\bar{\gamma}_{\omega}} \bar{e}_{\omega,t}. \quad (\text{A.1.4})$$

Since

$$\text{Var}(\bar{e}_{\omega,t}) = \sum_{i=1}^N \omega_i^2 \left( \sum_{l=0}^{\infty} a_{il}^2 \right) \leq \sigma^{*2} \sum_{i=1}^N \omega_i^2 = O\left(\frac{1}{N}\right), \quad (\text{A.1.5})$$

it follows that as  $N \rightarrow \infty$ ,  $\bar{e}_{\omega,t} \rightarrow 0$ . Consequently the factor  $f_t$  can now be approximated by:

$$f_t \approx \frac{1}{\bar{\gamma}_{\omega}} \Delta \bar{y}_{\omega,t} - \frac{1}{\bar{\gamma}_{\omega}} b \overline{y_{\omega,t-1}^3}. \quad (\text{A.1.6})$$

### A.2 Theorem 1

Let

$$\begin{aligned} \Delta y_{it} &= a_i + b_i y_{i,t-1}^3 + c_i \Delta \bar{y}_t + d_i \overline{y_{i,t-1}^3} + e_{it} \\ &= b_i y_{i,t-1}^3 + \gamma_i' x_t + e_{it}, \end{aligned} \quad (\text{A.2.1})$$

where

$$\begin{aligned} \gamma_i &\equiv (a_i, c_i, d_i)', \\ x_t &\equiv \left( 1, \Delta \bar{y}_t, \overline{y_{t-1}^3} \right)'. \end{aligned}$$

Then, in matrix formulation, (A.2.1) is

$$\Delta y_i = b_i y_{i,-1}^3 + X \gamma_i + e_i,$$

where

$$\begin{aligned} \Delta y_i &\equiv (\Delta y_{i1}, \Delta y_{i2}, \dots, \Delta y_{iT})', \\ X &\equiv \left( \tau, \overline{\Delta y}, \overline{y_{-1}^3} \right), \\ \tau &\equiv (1, 1, \dots, 1)'. \end{aligned}$$

The  $t$ -statistic for testing that  $b_i = 0$  is

$$t_i(N, T) = T^{1/2} \frac{y_{i,-1}^3 M \Delta y_i}{(\Delta y_i' M \Delta y_i)^{1/2} (y_{i,-1}^3 M y_{i,-1}^3)^{1/2}}, \quad (\text{A.2.2})$$

where

$$M \equiv I_T - X (X' X)^{-1} X'.$$

To find an expression of (A.2.2) in terms of nuisance parameters, we may assume that

$$\Delta y_{it} = \gamma_i f_t + \varepsilon_{it}, \quad (\text{A.2.3})$$

where  $\varepsilon_{it}$  are *iid*  $(0, \sigma^2)$ . In matrix formulation, this is

$$\Delta y_i = f \gamma_i + \varepsilon_i, \quad (\text{A.2.4})$$

where

$$f \equiv (f_1, f_2, \dots, f_T)'$$

As in Pesaran [2007], with  $\xi_i \equiv \varepsilon_i - \delta_i \bar{\varepsilon}$ ,  $\delta_i \equiv \gamma_i / \bar{\gamma}$ ,

$$\begin{aligned} \Delta y_i &= f \gamma_i + \varepsilon_i = \left( \frac{\gamma_i}{\bar{\gamma}} \bar{\gamma} f + \frac{\gamma_i}{\bar{\gamma}} \bar{\varepsilon} \right) + \left( \varepsilon_i - \frac{\gamma_i}{\bar{\gamma}} \bar{\varepsilon} \right) \\ &= \delta_i \overline{\Delta y} + \xi_i, \end{aligned}$$

where

$$\xi_i \equiv \varepsilon_i - \delta_i \bar{\varepsilon}.$$

Hence, because  $M \overline{\Delta y} = 0$ ,

$$M \Delta y_i = M_i \xi_i = \omega_i M v_i, \quad (\text{A.2.5})$$

where  $\omega_i^2 = \text{Var}(\xi_i)$  and  $v_i$  is standard normal. Similarly, with  $y_0 = 0$ ,

$$\begin{aligned} y_{i,t-1} &= \gamma_i s_{f,t-1} + s_{i,t-1} \\ &= \left( \frac{\gamma_i}{\bar{\gamma}} \bar{\gamma} s_{f,t-1} + \frac{\gamma_i}{\bar{\gamma}} \bar{s}_{t-1} \right) + \left( s_{i,t-1} - \frac{\gamma_i}{\bar{\gamma}} \bar{s}_{t-1} \right) \\ &= \delta_i \bar{y}_{t-1} + \eta_{i,t-1}, \end{aligned}$$

where

$$\eta_{i,t} \equiv s_{i,t} - \delta_i \bar{s}_t.$$

The corresponding expression for  $y_{i,t-1}^3$  is

$$\begin{aligned} y_{i,t-1}^3 &= (\delta_i \bar{y}_{t-1} + \eta_{i,t-1})^3 \\ &= \delta_i^3 \bar{y}_{t-1}^3 + 3\delta_i^2 \bar{y}_{t-1}^2 \eta_{i,t-1} + 3\delta_i \bar{y}_{t-1} \eta_{i,t-1}^2 + \eta_{i,t-1}^3. \end{aligned} \quad (\text{A.2.6})$$

Hence, stacking the  $\bar{y}_{i,t-1}^j$  in vectors  $\bar{y}_{t-1}^j$ , for  $j = 1, 2, 3$ , we have, because  $M\bar{y}_{-1}^3 = 0$ ,

$$My_{i,-1}^3 = 3\delta_i^2 M(\bar{y}_{-1}^2 \odot \eta_{i,-1}) + 3\delta_i M(\bar{y}_{-1} \odot \eta_{i,-1}^2) + M\eta_{i,-1}^3,$$

where  $\odot$  is the Hadamard (elementwise) product. Moreover, because

$$\bar{y}_{t-1} = \bar{\gamma} s_{f,t-1} + \bar{s}_{t-1}, \quad (\text{A.2.7})$$

the  $t$ th element of  $\bar{y}_{-1}^2$  is

$$\bar{y}_{t-1}^2 = \bar{\gamma}^2 s_{f,t-1}^2 + 2\bar{\gamma} s_{f,t-1} \bar{s}_{t-1} + \bar{s}_{t-1}^2,$$

and it follows that

$$\begin{aligned} My_{i,-1}^3 &= 3\delta_i^2 \bar{\gamma}^2 M(s_{f,-1}^2 \odot \eta_{i,-1}) + 6\delta_i^2 \bar{\gamma} M(s_{f,-1} \odot \bar{s}_{-1} \odot \eta_{i,-1}) \\ &\quad + 3\delta_i^2 M(\bar{s}_{-1}^2 \odot \eta_{i,-1}) + 3\delta_i \bar{\gamma} M(s_{f,-1} \odot \eta_{i,-1}^2) \\ &\quad + 3\delta_i M(\bar{s}_{-1} \odot \eta_{i,-1}^2) + M\eta_{i,-1}^3 \\ &= \omega_i^3 M\mathfrak{s}_{i,-1}^3 + R_{i1} + R_{i2}, \end{aligned} \quad (\text{A.2.8})$$

where

$$\begin{aligned} R_{i1} &\equiv 3\gamma_i^2 \omega_i M(s_{f,-1}^2 \odot \mathfrak{s}_{i,-1}) + 3\gamma_i \omega_i^2 M(s_{f,-1} \odot \mathfrak{s}_{i,-1}^2), \\ R_{i2} &\equiv 6\gamma_i^2 \bar{\gamma}^{-1} \omega_i M(s_{f,-1} \odot \bar{s}_{-1} \odot \mathfrak{s}_{i,-1}) + 3\gamma_i^2 \bar{\gamma}^{-2} \omega_i M(\bar{s}_{-1}^2 \odot \mathfrak{s}_{i,-1}) \\ &\quad + 3\gamma_i \bar{\gamma}^{-1} \omega_i^2 M(\bar{s}_{-1} \odot \mathfrak{s}_{i,-1}^2) \end{aligned}$$

with  $\mathfrak{s}_{i,-1} = \eta_{i,-1}/\omega_i$ . Hence, via [Equation A.2.5](#), the nominator of [Equation A.2.2](#) is

$$y_{i,-1}^3 M \Delta y_i = \omega_i^4 v_i' M \mathfrak{s}_{i,-1}^3 + \omega_i v_i' R_{i1} + \omega_i v_i' R_{i2} \quad (\text{A.2.9})$$

Similarly, via [Equation A.2.8](#),

$$\begin{aligned} y_{i,-1}^3 M y_{i,-1}^3 &= (M y_{i,-1}^3)' M y_{i,-1}^3 \\ &= \omega_i^6 \mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3 + \omega_i^3 \mathfrak{s}_{i,-1}^{3'} M R_{i1} + \omega_i^3 R_{i1}' M \mathfrak{s}_{i,-1}^3 \\ &\quad + \omega_i^3 \mathfrak{s}_{i,-1}^{3'} M R_{i2} + \omega_i^3 R_{i2}' M \mathfrak{s}_{i,-1}^3 \\ &\quad + (R_{i1} + R_{i2})' (R_{i1} + R_{i2}). \end{aligned} \quad (\text{A.2.10})$$

As in [Pesaran \[2007\]](#), the terms involving  $\bar{s}_{-1}$ , which are collected in  $R_{i2}$ , will tend to zero as  $N \rightarrow \infty$ . Hence, as  $N \rightarrow \infty$ , using [Equations Equation A.2.5-Equation A.2.10](#), [Equation A.2.2](#) is asymptotically equivalent to

$$\begin{aligned} &t_i^*(N, T) \\ &\equiv T^{1/2} \frac{\omega_i^4 v_i' M \mathfrak{s}_{i,-1}^3 + \omega_i v_i' R_{i1}}{\sqrt{\omega_i^2 v_i' M v_i} \sqrt{\omega_i^6 \mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3 + \omega_i^3 (\mathfrak{s}_{i,-1}^{3'} M R_{i1} + R_{i1}' M \mathfrak{s}_{i,-1}^3) + R_{i1}' R_{i1}}} \\ &= T^{1/2} \frac{v_i' M \mathfrak{s}_{i,-1}^3 + \omega_i^{-3} v_i' R_{i1}}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3 + \omega_i^{-3} (\mathfrak{s}_{i,-1}^{3'} M R_{i1} + R_{i1}' M \mathfrak{s}_{i,-1}^3) + \omega_i^{-6} R_{i1}' R_{i1}}}, \end{aligned}$$

and by Taylor expansion,

$$t_i^*(N, T) = T^{1/2} \frac{v_i' M \mathfrak{s}_{i,-1}^3}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} (1 + R_i),$$

where

$$R_i \equiv \frac{\omega_i^{-3} v_i' R_{i1}}{v_i' M \mathfrak{s}_{i,-1}^3} - \frac{1}{2} \frac{\omega_i^{-3} (\mathfrak{s}_{i,-1}^{3'} M R_{i1} + R_{i1}' M \mathfrak{s}_{i,-1}^3)}{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3} + O(\|R_{i1}^2\|).$$

Now, as  $N \rightarrow \infty$ , the mean statistic

$$\bar{t}(N, T) \equiv \frac{1}{N} \sum_{i=1}^N t_i(N, T),$$

is asymptotically equivalent to

$$\bar{t}^*(N, T) \equiv \frac{1}{N} \sum_{i=1}^N t_i^*(N, T) = \frac{T^{1/2}}{N} \sum_{i=1}^N \frac{v_i' M \mathfrak{s}_{i,-1}^3}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} + R, \quad (\text{A.2.11})$$

where

$$R \equiv \frac{T^{1/2}}{N} \sum_{i=1}^N \frac{v_i' M \mathfrak{s}_{i,-1}^3}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} \left\{ \frac{\omega_i^{-3} v_i' R_{i1}}{v_i' M \mathfrak{s}_{i,-1}^3} - \frac{1}{2} \frac{\omega_i^{-3} (\mathfrak{s}_{i,-1}^{3'} M R_{i1} + R_{i1}' M \mathfrak{s}_{i,-1}^3)}{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3} + O(\|R_{i1}^2\|) \right\}. \quad (\text{A.2.12})$$

Here,

$$\frac{T^{1/2}}{N} \sum_{i=1}^N \frac{v_i' M \mathfrak{s}_{i,-1}^3}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} \frac{\omega_i^{-3} v_i' R_{i1}}{v_i' M \mathfrak{s}_{i,-1}^3} = \frac{3\omega_i^{-2}}{N} \sum_{i=1}^N \gamma_i \kappa_i,$$

where

$$\kappa_i \equiv T^{1/2} \frac{v_i' \{ \gamma_i M (s_{f,-1}^2 \odot \mathfrak{s}_{i,-1}) + \omega_i M (s_{f,-1} \odot \mathfrak{s}_{i,-1}^2) \}}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}},$$

which is  $O_p(1)$ . Now, by the Cauchy-Schwarz inequality,

$$\begin{aligned} & \left( \frac{T^{1/2}}{N} \sum_{i=1}^N \frac{v_i' M \mathfrak{s}_{i,-1}^3}{\sqrt{v_i' M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} \frac{\omega_i^{-3} v_i' R_{i1}}{v_i' M \mathfrak{s}_{i,-1}^3} \right)^2 \\ & \leq (3\omega_i^{-2})^2 \left( \frac{1}{N} \sum_{i=1}^N \kappa_i^2 \right) \left( \frac{1}{N} \sum_{i=1}^N \gamma_i^2 \right) \\ & = O_p(1) \left( \frac{1}{N} \sum_{i=1}^N \gamma_i^2 \right), \end{aligned}$$

which tends to zero as  $N \rightarrow \infty$  if

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{i=1}^N \gamma_i^2 = 0,$$

which is Assumption 7. The same argument may be applied to all terms composing [Equation A.2.12](#), and so, [Equation A.2.11](#) implies that as  $N \rightarrow \infty$ ,  $\bar{t}(N, T)$  is asymptotically equivalent to

$$\frac{T^{1/2}}{N} \sum_{i=1}^N \frac{v'_i M \mathfrak{s}_{i,-1}^3}{\sqrt{v'_i M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}}.$$

Following the lines of proof as in [Pesaran \[2007\]](#), we have

$$T^{-2} v'_i M \mathfrak{s}_{i,-1}^3 = T^{-2} v'_i \mathfrak{s}_{i,-1}^3 - (v'_i X D) (D X' X D)^{-1} (T^{-2} D X' \mathfrak{s}_{i,-1}^3),$$

where

$$D = \begin{pmatrix} T^{-1/2} & 0 & 0 \\ 0 & T^{-1/2} & 0 \\ 0 & 0 & T^{-2} \end{pmatrix}.$$

Moreover,

$$\begin{aligned} D X' v_i &= \begin{pmatrix} T^{-1/2} \tau' v_i \\ T^{-1/2} \overline{\Delta y}' v_i \\ T^{-2} \overline{y}_-^3 v_i \end{pmatrix}, \\ T^{-2} D X' \mathfrak{s}_{i,-1}^3 &= \begin{pmatrix} T^{-5/2} \tau' \mathfrak{s}_{i,-1}^3 \\ T^{-5/2} \overline{\Delta y}' \mathfrak{s}_{i,-1}^3 \\ T^{-4} \overline{y}_-^3 \mathfrak{s}_{i,-1}^3 \end{pmatrix}, \\ D X' X D &= \begin{pmatrix} 1 & T^{-1} \tau' \overline{\Delta y} & T^{-5/2} \tau' \overline{y}_-^3 \\ T^{-1} \tau' \overline{\Delta y} & T^{-1} \overline{\Delta y}' \overline{\Delta y} & T^{-5/2} \overline{\Delta y}' \overline{y}_-^3 \\ T^{-5/2} \tau' \overline{y}_-^3 & T^{-5/2} \overline{\Delta y}' \overline{y}_-^3 & T^{-4} \overline{y}_-^3 \overline{y}_-^3 \end{pmatrix}, \end{aligned}$$

where, because from [Equation A.2.4](#) and [Equation A.2.6](#), with obvious notation,

$$\begin{aligned} \overline{\Delta y} &= \bar{\gamma} f + \bar{\varepsilon} \\ \overline{y}_-^3 &= \bar{\delta}^3 \overline{y}_{-1}^3 + 3(\bar{\delta}^2 \odot \eta_{-1}) \overline{y}_{-1}^2 + 3(\bar{\delta} \odot \eta_{-1}^2) \overline{y}_{-1} + \bar{\eta}_{-1}^3 \end{aligned}$$

we have

$$\begin{aligned} T^{-1} \tau' \overline{\Delta y} &= \bar{\gamma} T^{-1} \tau' f + T^{-1} \tau' \bar{\varepsilon}, \\ T^{-1} \overline{\Delta y}' \overline{\Delta y} &= \bar{\gamma}^2 T^{-1} f' f + 2\bar{\gamma} T^{-1} f' \bar{\varepsilon} + T^{-1} \bar{\varepsilon}' \bar{\varepsilon}, \\ T^{-5/2} \tau' \overline{y}_-^3 &= \bar{\delta}^3 T^{-5/2} \tau' \overline{y}_{-1}^3 + 3(\bar{\delta}^2 \odot \eta_{-1}) T^{-5/2} \tau' \overline{y}_{-1}^2 \\ &\quad + 3(\bar{\delta} \odot \eta_{-1}^2) T^{-5/2} \tau' \overline{y}_{-1} + T^{-5/2} \tau' \bar{\eta}_{-1}^3, \\ T^{-5/2} \overline{\Delta y}' \overline{y}_-^3 &= \bar{\gamma} \bar{\delta}^3 T^{-5/2} f' \overline{y}_{-1}^3 + 3\bar{\gamma} (\bar{\delta}^2 \odot \eta_{-1}) T^{-5/2} f' \overline{y}_{-1}^2 \\ &\quad + 3\bar{\gamma} (\bar{\delta} \odot \eta_{-1}^2) T^{-5/2} f' \overline{y}_{-1} + \bar{\gamma} T^{-5/2} f' \bar{\eta}_{-1}^3 \\ &\quad + \bar{\delta}^3 T^{-5/2} \bar{\varepsilon}' \overline{y}_{-1}^3 + 3(\bar{\delta}^2 \odot \eta_{-1}) T^{-5/2} \bar{\varepsilon}' \overline{y}_{-1}^2 \\ &\quad + 3(\bar{\delta} \odot \eta_{-1}^2) T^{-5/2} \bar{\varepsilon}' \overline{y}_{-1} + T^{-5/2} \bar{\varepsilon}' \bar{\eta}_{-1}^3, \\ T^{-4} \overline{y}_-^3 \overline{y}_-^3 &= T^{-4} \bar{\delta}^3 \overline{y}_{-1}^3 \overline{y}_{-1}^3 + 6\bar{\delta}^3 (\bar{\delta}^2 \odot \eta_{-1}) T^{-4} \overline{y}_{-1}^3 \overline{y}_{-1}^2 \\ &\quad + 6\bar{\delta}^3 (\bar{\delta} \odot \eta_{-1}^2) T^{-4} \overline{y}_{-1}^3 \overline{y}_{-1} + 2\bar{\delta}^3 T^{-4} \overline{y}_{-1}^3 \bar{\eta}_{-1}^3 \\ &\quad + 9(\bar{\delta}^2 \odot \eta_{-1})^2 T^{-4} \overline{y}_{-1}^2 \overline{y}_{-1}^2 + 18(\bar{\delta}^2 \odot \eta_{-1}) (\bar{\delta} \odot \eta_{-1}^2) T^{-4} \overline{y}_{-1}^2 \overline{y}_{-1} \\ &\quad + 6(\bar{\delta}^2 \odot \eta_{-1}) T^{-4} \overline{y}_{-1}^2 \bar{\eta}_{-1}^3 + 9(\bar{\delta} \odot \eta_{-1}^2)^2 T^{-4} \overline{y}_{-1} \overline{y}_{-1} \\ &\quad + 6(\bar{\delta} \odot \eta_{-1}^2) T^{-4} \overline{y}_{-1} \bar{\eta}_{-1}^3 + T^{-4} \bar{\eta}_{-1}^3 \bar{\eta}_{-1}^3. \end{aligned}$$

As in [Pesaran \[2007\]](#), as  $N \rightarrow \infty$ , terms involving  $\bar{\varepsilon}$ ,  $\overline{\eta^3}$  or  $\bar{s}_-$  may be neglected. In particular, it follows from [\(A.2.7\)](#) that  $\bar{y}_{-1}$  is dominated by  $\bar{\gamma}s_{f,-1}$ . Hence, denoting asymptotic equivalence by  $\sim$ , and cancelling out lower order terms we have, because  $\mathbf{s}_{i,-1} \sim \sigma_i^{-1}s_{i,-1}$ ,

$$T^{-2}v'_i\mathbf{s}_{i,-1}^3 \sim \sigma_i^{-3}T^{-2}\varepsilon'_i s_{i,-1}^3,$$

and moreover,

$$\begin{aligned} T^{-1}\tau'\overline{\Delta y} &\sim \bar{\gamma}T^{-1}\tau'f, \\ T^{-1}\overline{\Delta y}'\overline{\Delta y} &\sim \bar{\gamma}^2T^{-1}f'f, \\ T^{-5/2}\tau'\overline{y_-^3} &\sim \bar{\gamma}^3\bar{\delta}^3T^{-5/2}\tau's_{f,-1}^3, \\ T^{-5/2}\overline{\Delta y}'\overline{y_-^3} &\sim \bar{\gamma}^4\bar{\delta}^3T^{-5/2}f's_{f,-1}^3, \\ T^{-4}\overline{y_-^3}'\overline{y_-^3} &\sim \bar{\gamma}^6\bar{\delta}^3T^{-4}s_{f,-1}^3s_{f,-1}^3, \end{aligned}$$

and so, denoting the limits of  $\bar{\gamma}$  and  $\bar{\delta}^3$  by  $\gamma_*$  and  $\delta_*$  (from now on,  $\rightarrow$  denotes convergence in distribution),

$$DX'XD \rightarrow \Gamma\tilde{\Psi}_T\Gamma,$$

where

$$\Gamma \equiv \begin{pmatrix} 1 & 0 & 0 \\ 0 & \gamma_* & 0 \\ 0 & 0 & \gamma_*^3\delta_* \end{pmatrix},$$

$$\tilde{\Psi}_T \equiv \begin{pmatrix} 1 & T^{-1}\tau'f & T^{-5/2}\tau's_{f,-1}^3 \\ T^{-1}\tau'f & T^{-1}f'f & T^{-5/2}f's_{f,-1}^3 \\ T^{-5/2}\tau's_{f,-1}^3 & T^{-5/2}f's_{f,-1}^3 & T^{-4}s_{f,-1}^3s_{f,-1}^3 \end{pmatrix}.$$

Similarly, because  $v_i \sim \sigma_i^{-1}\varepsilon_i$ ,

$$\begin{aligned} DXv_i &\rightarrow \sigma_i^{-1}\Gamma\tilde{q}_T, \\ T^{-2}DX'\mathbf{s}_{i,-1}^3 &\rightarrow \sigma_i^{-3}\Gamma\tilde{h}_T, \end{aligned}$$

where

$$\tilde{q}_T = \begin{pmatrix} T^{-1/2}\tau'\varepsilon_i \\ T^{-1/2}f'\varepsilon_i \\ T^{-2}s_{f,-1}^3\varepsilon_i \end{pmatrix}, \quad \tilde{h}_T = \begin{pmatrix} T^{-5/2}\tau's_{i,-1}^3 \\ T^{-5/2}f's_{i,-1}^3 \\ T^{-4}s_{f,-1}^3s_{i,-1}^3 \end{pmatrix},$$

and so, as  $N \rightarrow \infty$ ,

$$T^{-2}v'_iM\mathbf{s}_{i,-1}^3 \rightarrow \sigma_i^{-4} \left( T^{-2}\varepsilon'_i s_{i,-1}^3 - \tilde{q}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T \right).$$

In the same fashion,

$$T^{-1}v'_iMv_i \rightarrow \sigma_i^{-2} \left( T^{-1}\varepsilon'_i\varepsilon_i - T^{-1}\tilde{q}'_T \tilde{\Psi}_T^{-1} \tilde{q}_T \right),$$

and

$$T^{-4}\mathbf{s}_{i,-1}^3M\mathbf{s}_{i,-1}^3 \rightarrow \sigma_i^{-6} \left( T^{-4}s_{i,-1}^3s_{i,-1}^3 - \tilde{h}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T \right),$$

implying that for all  $i$ , as  $N \rightarrow \infty$ ,

$$\begin{aligned} & T^{1/2} \frac{v'_i M \mathfrak{s}_{i,-1}^3}{\sqrt{v'_i M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} \\ \rightarrow & \frac{T^{-2} \varepsilon'_i \mathfrak{s}_{i,-1}^3 - \tilde{q}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T}{\sqrt{T^{-1} \varepsilon'_i \varepsilon_i - T^{-1} \tilde{q}'_T \tilde{\Psi}_T^{-1} \tilde{q}_T} \sqrt{T^{-4} \mathfrak{s}_{i,-1}^{3'} \mathfrak{s}_{i,-1}^3 - \tilde{h}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T}}. \end{aligned}$$

Note that the r.h.s. of this expression does not depend on any nuisance parameters. This completes the proof of Theorem 1.

### A.3 Theorem 2

Without loss of generality, set  $\sigma_i = 1$ . As in Pesaran [2007], as  $T \rightarrow \infty$  (we use shorthand notation for the integrals so that e.g.  $\int W_i^3 dW_i = \int_0^1 W_i(t)^3 dW_i(t)$  and  $\int W_i^6 = \int_0^1 W_i(t)^3 dt$ )

$$\begin{aligned} T^{-2} \varepsilon'_i \mathfrak{s}_{i,-1}^3 & \rightarrow \int W_i^3 dW_i, \\ T^{-1} \varepsilon'_i \varepsilon_i & \rightarrow 1, \\ T^{-4} \mathfrak{s}_{i,-1}^{3'} \mathfrak{s}_{i,-1}^3 & \rightarrow \int W_i^6, \\ \tilde{\Psi}_T & \rightarrow \begin{pmatrix} 1 & 0 & \int W_f^3 \\ 0 & 1 & 0 \\ \int W_f^3 & 0 & \int W_f^6 \end{pmatrix} \equiv \tilde{\Psi}, \\ \tilde{q}_T & \rightarrow \begin{pmatrix} W_i(1) \\ W_{fi}(1) \\ \int W_f^3 dW_i \end{pmatrix} \equiv \tilde{q}, \quad \tilde{h}_T \rightarrow \begin{pmatrix} \int W_i^3 \\ 0 \\ \int W_f^3 W_i^3 \end{pmatrix} \equiv \tilde{h}, \end{aligned}$$

where  $W_i, W_f$  and  $W_{fi}$  are standard Brownian motions. (Moreover,  $W_i$  and  $W_f$  are mutually independent.) Hence, it follows that  $T^{-1} q'_T \Psi_T^{-1} h_T$  may be neglected, and moreover,

$$\begin{aligned} \tilde{q}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T & \rightarrow \tilde{q}' \tilde{\Psi}^{-1} \tilde{h} = q' \Psi^{-1} h, \\ \tilde{h}'_T \tilde{\Psi}_T^{-1} \tilde{h}_T & \rightarrow \tilde{h}' \tilde{\Psi}^{-1} \tilde{h} = h' \Psi^{-1} h, \end{aligned}$$

where

$$\begin{aligned} \Psi & = \begin{pmatrix} 1 & \int W_f^3 \\ \int W_f^3 & \int W_f^6 \end{pmatrix}, \\ q & = \begin{pmatrix} W_i(1) \\ \int W_f^3 dW_i \end{pmatrix}, \quad h = \begin{pmatrix} \int W_i^3 \\ \int W_f^3 W_i^3 \end{pmatrix}, \end{aligned}$$

and we find the sequential limit result

$$T^{1/2} \frac{v'_i M \mathfrak{s}_{i,-1}^3}{\sqrt{v'_i M v_i} \sqrt{\mathfrak{s}_{i,-1}^{3'} M \mathfrak{s}_{i,-1}^3}} \rightarrow \frac{\int W_i^3 dW_i - q' \Psi^{-1} h}{\sqrt{\int W_i^6 - h' \Psi^{-1} h}}. \quad (\text{A.3.1})$$

Along the lines of Pesaran [2007], it may be proved that this is also the joint limit result as  $N, T \rightarrow \infty$  simultaneously such that  $N/T \rightarrow k$ , where  $k$  is a finite and positive constant.



## B Critical Values

### B.1 Individual NCADF Distribution

Table 11: Critical Values of Individual NCADF Distribution

N	T	1 %	2.5 %	5 %	10 %	N	T	1 %	2.5 %	5 %	10 %
10	10	-5.18	-4.17	-3.50	-2.87	50	10	-5.16	-4.17	-3.52	-2.91
	15	-4.19	-3.60	-3.16	-2.69		15	-4.21	-3.57	-3.15	-2.68
	20	-3.93	-3.44	-3.07	-2.67		20	-4.10	-3.47	-3.11	-2.69
	30	-3.79	-3.38	-3.05	-2.70		30	-3.75	-3.33	-3.00	-2.69
	50	-3.81	-3.41	-3.11	-2.78		50	-3.68	-3.35	-3.04	-2.76
	70	-3.67	-3.39	-3.12	-2.80		70	-3.70	-3.36	-3.07	-2.75
	100	-3.71	-3.39	-3.12	-2.80		100	-3.59	-3.31	-3.09	-2.79
	200	-3.73	-3.40	-3.12	-2.82		200	-3.72	-3.36	-3.10	-2.81
15	10	-5.35	-4.22	-3.52	-2.92	70	10	-5.17	-4.23	-3.52	-2.92
	15	-4.21	-3.64	-3.15	-2.67		15	-4.32	-3.64	-3.22	-2.74
	20	-3.96	-3.42	-3.06	-2.68		20	-3.97	-3.47	-3.10	-2.65
	30	-3.81	-3.36	-3.06	-2.69		30	-3.79	-3.41	-3.06	-2.71
	50	-3.69	-3.32	-3.06	-2.75		50	-3.73	-3.41	-3.11	-2.76
	70	-3.75	-3.41	-3.11	-2.78		70	-3.68	-3.37	-3.05	-2.76
	100	-3.70	-3.38	-3.13	-2.76		100	-3.71	-3.40	-3.10	-2.81
	200	-3.67	-3.37	-3.09	-2.78		200	-3.62	-3.34	-3.11	-2.83
20	10	-5.05	-4.20	-3.47	-2.89	100	10	-4.89	-3.99	-3.39	-2.81
	15	-4.27	-3.63	-3.13	-2.73		15	-4.04	-3.53	-3.16	-2.75
	20	-3.94	-3.39	-3.04	-2.67		20	-3.91	-3.45	-3.05	-2.66
	30	-3.71	-3.39	-3.09	-2.74		30	-3.76	-3.36	-3.06	-2.70
	50	-3.70	-3.28	-3.04	-2.73		50	-3.63	-3.33	-3.04	-2.75
	70	-3.66	-3.35	-3.07	-2.75		70	-3.64	-3.31	-3.01	-2.74
	100	-3.74	-3.38	-3.09	-2.80		100	-3.74	-3.35	-3.10	-2.79
	200	-3.77	-3.40	-3.14	-2.84		200	-3.69	-3.40	-3.11	-2.82
30	10	-5.62	-4.37	-3.55	-2.95	200	10	-5.21	-4.17	-3.42	-2.84
	15	-4.22	-3.62	-3.14	-2.68		15	-4.30	-3.67	-3.21	-2.78
	20	-3.87	-3.42	-3.09	-2.70		20	-3.91	-3.44	-3.11	-2.70
	30	-3.86	-3.42	-3.14	-2.73		30	-3.69	-3.34	-3.04	-2.73
	50	-3.69	-3.37	-3.06	-2.75		50	-3.77	-3.40	-3.10	-2.77
	70	-3.71	-3.32	-3.07	-2.75		70	-3.66	-3.28	-3.08	-2.75
	100	-3.77	-3.32	-3.10	-2.79		100	-3.70	-3.38	-3.11	-2.79
	200	-3.68	-3.37	-3.11	-2.84		200	-3.64	-3.38	-3.14	-2.81

## B.2 Panel NCADF Distribution

Table 12: Critical values of Panel NCADF Distribution

N	T	1%	2.5%	5%	10%	N	T	1%	2.5%	5%	10%
10	200	-2.50	-2.40	-2.33	-2.25	50	200	-2.14	-2.09	-2.04	-1.99
10	100	-2.42	-2.31	-2.22	-2.11	50	100	-2.10	-2.05	-2.01	-1.96
10	70	-2.39	-2.27	-2.19	-2.10	50	70	-2.08	-2.03	-1.99	-1.94
10	50	-2.36	-2.26	-2.16	-2.05	50	50	-2.05	-2.00	-1.96	-1.91
10	30	-2.31	-2.20	-2.12	-2.01	50	30	-2.00	-1.95	-1.90	-1.84
10	20	-2.32	-2.20	-2.09	-1.97	50	20	-1.96	-1.90	-1.85	-1.79
10	15	-2.34	-2.19	-2.08	-1.94	50	15	-1.95	-1.88	-1.82	-1.75
10	10	-2.53	-2.34	-2.17	-1.98	50	10	-2.01	-1.91	-1.83	-1.75
15	200	-2.33	-2.25	-2.18	-2.09	70	200	-2.11	-2.06	-2.02	-1.98
15	100	-2.30	-2.22	-2.14	-2.06	70	100	-2.07	-2.03	-1.99	-1.95
15	70	-2.26	-2.19	-2.13	-2.04	70	70	-2.05	-2.00	-1.97	-1.92
15	50	-2.24	-2.16	-2.08	-2.00	70	50	-2.02	-1.98	-1.94	-1.89
15	30	-2.20	-2.11	-2.03	-1.95	70	30	-1.96	-1.91	-1.87	-1.83
15	20	-2.17	-2.09	-2.00	-1.90	70	20	-1.92	-1.87	-1.83	-1.77
15	15	-2.19	-2.08	-1.98	-1.88	70	15	-1.91	-1.84	-1.80	-1.73
15	10	-2.34	-2.18	-2.04	-1.90	70	10	-1.95	-1.88	-1.80	-1.72
20	200	-2.26	-2.19	-2.13	-2.06	100	200	-2.08	-2.04	-2.01	-1.97
20	100	-2.24	-2.16	-2.11	-2.03	100	100	-2.05	-2.01	-1.97	-1.93
20	70	-2.20	-2.13	-2.08	-2.00	100	70	-2.02	-1.99	-1.95	-1.91
20	50	-2.18	-2.11	-2.05	-1.98	100	50	-1.99	-1.95	-1.92	-1.88
20	30	-2.14	-2.07	-2.00	-1.92	100	30	-1.94	-1.89	-1.86	-1.81
20	20	-2.11	-2.03	-1.95	-1.86	100	20	-1.89	-1.84	-1.81	-1.76
20	15	-2.10	-2.00	-1.93	-1.84	100	15	-1.87	-1.82	-1.77	-1.72
20	10	-2.22	-2.09	-1.97	-1.84	100	10	-1.92	-1.85	-1.78	-1.70
30	200	-2.20	-2.14	-2.09	-2.02	200	200	-2.05	-2.01	-1.99	-1.95
30	100	-2.18	-2.11	-2.06	-2.00	200	100	-2.01	-1.98	-1.96	-1.92
30	70	-2.15	-2.09	-2.03	-1.97	200	70	-2.00	-1.96	-1.93	-1.89
30	50	-2.11	-2.05	-2.00	-1.94	200	50	-1.96	-1.93	-1.90	-1.86
30	30	-2.07	-2.00	-1.95	-1.88	200	30	-1.90	-1.87	-1.84	-1.80
30	20	-2.02	-1.95	-1.90	-1.83	200	20	-1.86	-1.81	-1.78	-1.73
30	15	-2.02	-1.94	-1.87	-1.79	200	15	-1.82	-1.78	-1.74	-1.69
30	10	-2.13	-2.00	-1.90	-1.80	200	10	-1.87	-1.80	-1.75	-1.68

## C Tables when Assumption 7 is relaxed

Table 13: Size of Cross Sectionally Augmented Nonlinear Panel Unit Root Tests  
No Serial Correlation, Weak and Strong Cross Section Dependence Case

		Weak Cross Section Dependence					Strong Cross Section Dependence				
		$T$					$T$				
$N$	Test	10	20	30	50	100	10	20	30	50	100
10	CIPS	.0498	.0492	.0540	.0506	.0496	.0550	.0492	.0594	.0520	.0616
	NCIPS	.0468	.0474	.0582	.0438	.0494	.0508	.0432	.0456	.0414	.0474
20	CIPS	.0538	.0508	.0464	.0520	.0520	.0488	.0492	.0566	.0568	.0614
	NCIPS	.0532	.0484	.0444	.0556	.0488	.0468	.0432	.0448	.0402	.0432
30	CIPS	.0554	.0560	.0426	.0498	.0490	.0568	.0518	.0568	.0466	.0504
	NCIPS	.0516	.0456	.0490	.0478	.0448	.0470	.0394	.0438	.0332	.0354
50	CIPS	.0516	.0564	.0508	.0432	.0496	.0616	.0566	.0422	.0448	.0458
	NCIPS	.0474	.0520	.0486	.0474	.0512	.0408	.0410	.0386	.0328	.0348
100	CIPS	.0526	.0454	.0490	.0468	.0488	.0522	.0518	.0496	.0530	.0500
	NCIPS	.0470	.0458	.0434	.0452	.0478	.0436	.0414	.0348	.0354	.0350

Table 14: Power of Cross Sectionally Augmented Nonlinear Panel Unit Root Tests  
No Serial Correlation, Weak and Strong Cross Section Dependence Case  
Weak Alternative

		Weak Cross Section Dependence					Strong Cross Section Dependence				
		$T$					$T$				
$N$	Test	10	20	30	50	100	10	20	30	50	100
10	CIPS	.0584	.0884	.1356	.2398	.7280	.0632	.1144	.2294	.4952	.9180
	NCIPS	.0598	.1098	.1934	.3620	.8942	.0680	.1590	.3314	.6666	.9722
20	CIPS	.0590	.1134	.1594	.3960	.9480	.0554	.1278	.2678	.7026	.9918
	NCIPS	.0818	.1724	.2970	.6516	.9954	.0820	.2240	.4618	.8780	.9986
30	CIPS	.0712	.1116	.1818	.4696	.9864	.0486	.1302	.3234	.8080	.9964
	NCIPS	.0906	.2002	.3806	.7988	.9998	.0842	.2508	.5628	.9536	.9998
50	CIPS	.0660	.1246	.2088	.5280	1.000	.0516	.1466	.3638	.8916	1.000
	NCIPS	.0978	.2678	.5032	.9334	.0512	.0846	.3134	.6700	.9900	1.000
100	CIPS	.0744	.1264	.2398	.6428	1.000	.0478	.1476	.4122	.9592	1.000
	NCIPS	.1006	.3346	.6682	.9912	1.000	.0986	.3598	.7776	.9976	1.000

Table 15: Power of Cross Sectionally Augmented Nonlinear Panel Unit Root Tests  
 No Serial Correlation, Weak and Strong Cross Section Dependence Case  
 Strong Alternative

		Weak Cross Section Dependence					Strong Cross Section Dependence				
		$T$					$T$				
$N$	Test	10	20	30	50	100	10	20	30	50	100
10	CIPS	.0862	.1928	.4066	.8394	1.000	.0886	.3448	.7300	.9516	.9914
	NCIPS	.0994	.3190	.6336	.9616	1.000	.1378	.5020	.8490	.9846	.9986
20	CIPS	.0890	.2692	.5978	.9870	1.000	.0898	.4930	.8976	.9956	1.000
	NCIPS	.1582	.5706	.8988	.9998	1.000	.1958	.7346	.9656	.9982	1.000
30	CIPS	.1010	.3014	.6978	.9974	1.000	.0978	.5900	.9654	.9996	.9998
	NCIPS	.1722	.6862	.9716	1.000	1.000	.2268	.8488	.9948	1.000	1.000
50	CIPS	.1016	.4010	.8064	1.000	1.000	.1048	.7074	.9908	1.000	1.000
	NCIPS	.2158	.8592	.9984	1.000	1.000	.2752	.9250	.9990	1.000	1.000
100	CIPS	.1124	.3994	.9210	1.000	1.000	.1004	.7626	.9996	1.000	1.000
	NCIPS	.2596	.9700	1.000	1.000	1.000	.3150	.9720	1.000	1.000	1.000

Table 16: Size of Cross Sectionally Augmented Nonlinear Panel Unit Root Tests  
 High Cross Section Dependence Case

		Positive Serial Correlation					Negative Serial Correlation				
		$T$					$T$				
$N$	Test	10	20	30	50	100	10	20	30	50	100
10	CIPS	.5504	.1402	.1066	.0764	.0668	.5688	.1606	.1162	.0756	.0694
	NCIPS	.2928	.0600	.0646	.0514	.0534	.2544	.0396	.0326	.0294	.0384
20	CIPS	.6520	.1492	.1034	.0796	.0800	.6960	.1722	.1124	.0886	.0744
	NCIPS	.3534	.0610	.0558	.0542	.0526	.3038	.0262	.0228	.0306	.0306
30	CIPS	.6990	.1504	.0976	.0768	.0664	.7598	.1888	.1160	.0866	.0606
	NCIPS	.3774	.0534	.0478	.0528	.0484	.3194	.0244	.0238	.0240	.0306
50	CIPS	.7700	.1666	.0932	.0734	.0618	.8132	.2000	.1172	.0794	.0680
	NCIPS	.4174	.0482	.0458	.0538	.0374	.3620	.0216	.0146	.0228	.0300
100	CIPS	.8292	.1502	.1012	.0726	.0648	.8758	.2100	.1284	.0898	.0742
	NCIPS	.4672	.0476	.0464	.0448	.0432	.3896	.0164	.0142	.0208	.0240

Table 17: Power of Cross Sectionally Augmented Nonlinear Panel Unit Root Tests  
Strong alternative, High Cross Section Dependence Case

		Positive Serial Correlation					Negative Serial Correlation				
		$T$					$T$				
$N$	Test	10	20	30	50	100	10	20	30	50	100
10	CIPS	.5836	.2762	.4712	.8246	.9808	.5746	.3378	.5846	.8714	.9756
	NCIPS	.3840	.3684	.6998	.9592	.9978	.3004	.2394	.5534	.8966	.9898
20	CIPS	.6922	.3472	.6214	.9576	.9984	.6870	.4642	.7680	.9802	.9952
	NCIPS	.4968	.5556	.8986	.9966	.9998	.3814	.3466	.7468	.9846	.9990
30	CIPS	.7464	.3858	.7222	.9884	.9998	.7676	.5350	.8618	.9966	.9996
	NCIPS	.5470	.6476	.9580	.9994	1.000	.4356	.4034	.8466	.9962	1.000
50	CIPS	.8016	.4738	.8238	.9996	1.000	.8202	.6054	.9288	.9998	1.000
	NCIPS	.6320	.7670	.9916	1.000	1.000	.4936	.5024	.9230	.9996	1.000
100	CIPS	.8734	.4850	.9234	1.000	1.000	.8810	.6742	.9806	1.000	1.000
	NCIPS	.7290	.8826	.9984	1.000	1.000	.5552	.6038	.9688	1.000	1.000

## References

- N. S. Balke and T. B. Fomby. Threshold cointegration. *International Economic Review*, 38:627–645, 1997. [6](#)
- J. Breitung and H. M. Pesaran. Unit roots and cointegration in panels. In L. Matyas and P. Sevestre, editors, *The Econometrics of Panel Data (Third Edition)*. Kluwer Academic Publishers, 2007. [1](#)
- W. A. Brock and C. H. Hommes. Heterogeneous beliefs and routes to chaos in a simple asset pricing model. *Journal of Economic Dynamics & Control*, 22:1235–1274, 1998. [3](#)
- M. Cerrato and N. Sarantis. A bootstrap panel unit root test under cross-sectional dependence, with an application to ppp. *Computational Statistics & Data Analysis*, 51:4028–4037, 2007a. [1](#)
- M. Cerrato and N. Sarantis. Does purchasing power parity hold in emerging markets? evidence from a panel of black market exchange rates. *International Journal of Finance & Economics*, 12:427–444, 2007b. [1](#)
- R. Chang and A. Velasco. A model of financial crises in emerging markets. *Quarterly Journal of Economics*, 116(2):489–517, May 2001. [2](#)
- M.-H. Chiang, C.-M. Kuan, and C.-H. Lo. Panel unit root test under smooth transition. National Cheng Kung University, Taiwan, October 2007. [5](#)
- I. Choi. Unit root tests for panel data. *Journal of International Money & Finance*, 20:249–272, 2001. [3](#)
- R. B. Davies. Hypothesis testing when a nuisance parameter is present under the alternative. *Biometrika*, 74:33–43, 1987. [9](#)

- P. De Grauwe and M. Grimaldi. Heterogeneity of agents, transactions costs and the exchange rate. *Journal of Economic Dynamics & Control*, 29:691–719, 2005. 3
- B. Dumas. Dynamic equilibrium and the real exchange rate in a spatially separated world. *Review of Financial Studies*, 5:153–180, 1992. 1
- S. N. Durlauf and P. A. Johnson. Multiple-regime and cross-country growth behaviour. *Journal of Applied Econometrics*, 10:365–384, 1995. 3
- J. D. Emerson and C. Kao. Testing for structural change in panel data: Gdp growth, consumption growth, and productivity growth. *Economics Bulletin*, 3(14):1–12, 2006. 1
- D. Fiaschi and A. M. Lavezzi. Nonlinear economic growth: Some theory and cross-country evidence. *Journal of Development Economics*, 84:271–290, 2007. 3
- O. Galor and D. N. Weil. Population, technology and growth: From the malthusian regime to the demographic transition and beyond. *American Economic Review*, 90:806–828, 2000. 2
- G. Goswami, M. Shrikhande, and L. Wu. A dynamic equilibrium model of real exchange rates with general transaction costs. mimeo, Fordham University, 2002. 1
- K. Hadri. Testing for stationarity in heterogeneous panel data. *The Econometric Journal*, 3:148–161, 2000. 3
- K. Im, H. Pesaran, and Y. Shin. Testing for unit roots in heterogeneous panels. *Journal of Econometrics*, 115:53–74, 2003. 5, 7
- O. Jeanne and P. Masson. Currency crises, sunspots and markov-switching regimes. *Journal of International Economics*, 50:327–350, 2000. 2
- G. Kapetanios, Y. Shin, and A. Snell. Testing for a unit root in the nonlinear star framework. *Journal of Econometrics*, 112:359–379, 2003. 3, 9, 10
- Z. Liu and T. Stengos. Non-linearities in cross-country growth regressions: A semiparametric approach. *Journal of Applied Econometrics*, 14:527–538, 1999. 3
- R. Luukkonen, P. Saikkonen, and T. TerSasvirta. Testing linearity against smooth transition autoregressive models. *Biometrika*, 75:491–499, 1988. 9
- T. Lux. Herd behaviour, bubbles and crashes. *Economic Journal*, 105:881–896, 1995. 3
- K. Matsuyama. Growing through cycles. *Econometrica*, 67:335–347, 1999. 2
- P. Michael, A. R. Nobay, and D. A. Peel. Transactions costs and nonlinear adjustment in real exchange rates: An empirical investigation. *Journal of Political Economy*, 105:862–879, 1997. 2, 6
- C. J. Murray and D. H. Papell. The purchasing power persistence paradigm. *Journal of International Economics*, 56:1–19, 2002. 5
- C. J. Murray and D. H. Papell. The purchasing power puzzle is worse than you think. *Empirical Economics*, 30(3):783–790, October 2005. 5

- S. Ng. A simple test for nonstationarity in mixed panels. *Journal of Business & Economic Statistics*, 26(1):113–127, January 2008. [4](#)
- P. G. J. O’Connell. Market frictions and real exchange rates’. *Journal of International Money & Finance*, 17:71–95, 1998. [1](#)
- R. F. Peretto. Industrial development, technological change, and long-run growth. *Journal of Development Economics*, 59:389–417, 1999. [2](#)
- M. H. Pesaran. Estimation and inference in large heterogeneous panels with a multifactor error structure. *Econometrica*, 74(4):967–1012, July 2006. [7](#), [9](#), [A.1](#)
- M. H. Pesaran. A simple panel unit root test in the presence of cross section dependence. *Journal of Applied Econometrics*, 22(2):265–312, 2007. [\(document\)](#), [3](#), [5](#), [10](#), [3.2](#), [4](#), [11](#), [4.2](#), [5](#), [5](#), [6](#), [A.2](#), [A.2](#), [A.2](#), [A.3](#), [A.3](#)
- M. H. Pesaran and S. Tosetti. Large panels with common factors and spatial correlations. *IZA Discussion Paper*, 3032, 2007. [7](#)
- E. E. Peters. *Fractal Market Analysis: Applying Chaos Theory to Investment & Economics*. John Wiley & Sons, 1994. [3](#)
- D. E. Rapach and M. E. Wohar. Nonlinear models of real exchange rate behaviour: A re-examination. Mimeo, Seattle University, 2003. [2](#)
- N. Sarantis. Modelling nonlinearities in effective exchange rates. *Journal of International Money & Finance*, 18:27–45, 1999. [2](#)
- P. Sercu and C. Uppal, R. Van Hulle. The exchange rate in the presence of transaction costs: Implications for tests of the purchasing power parity. *Journal of Finance*, 50:1309–1319, 1995. [1](#)
- R. Sollis, S. Leybourne, and P. Newbold. Tests for symmetric and asymmetric nonlinear mean reversion in real exchange rates. *Journal of Money, Credit and Banking*, 34:686–700, 2002. [3](#)
- M. P. Taylor, D. A. Peel, and L. Sarno. Nonlinear mean reversion in real exchange rates: Toward a solution to the purchasing power parity puzzle. *International Economic Review*, 42:1015–1042, 2001. [2](#)
- F. Zilibotti. A rostovian model of endogenous growth and underdevelopment traps. *European Economic Review*, 39:1569–1602, 1995. [2](#)